

FORM PTO-1390 (REV 11-2000)	U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 35-213
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (if known, see 37 C.F.R. 1.5) 09/913159 Unknown
INTERNATIONAL APPLICATION NO. PCT/EP99/09759	INTERNATIONAL FILING DATE 3 December 1999	PRIORITY DATE CLAIMED 12 February 1999

TITLE OF INVENTION

VIRAL VACCINE

APPLICANT(S) FOR DO/EO/US

SCHREIBER

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The U.S. has been elected by the expiration of 19 months from the priority date (Article 31).
5. A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☒ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 34
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has **NOT** expired.
 - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☒ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 To 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information. PTO-1449/ International Search Report/ Sequence Listing/ Front page of the PCT Publication/ Certification of Translation/Letter re: Sequence Listing

ST/PTO 10 AUG 2001


U.S. APPLICATION NO. 09/913159		ATTORNEY'S DOCKET NUMBER 35-213							
21. <input checked="" type="checkbox"/> The following fees are submitted:		CALCULATIONS PTO USE ONLY							
BASIC NATIONAL FEE (37 C.F.R. 1.492(a)(1)-(5)): -- Neither international preliminary examination fee (37 C.F.R. 1.482) nor international search fee (37 C.F.R. 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO\$1000.00 -- International preliminary examination fee (37 C.F.R. 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO.....\$860.00 -- International preliminary examination fee (37 C.F.R. 1.482) not paid to USPTO but international search fee (37 C.F.R. 1.445(a)(2)) paid to USPTO\$710.00 -- International preliminary examination fee (37 C.F.R. 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690.00 -- International preliminary examination fee (37 C.F.R. 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4).....\$100.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>		<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:10%; text-align: right;">\$</td> <td style="width:40%; text-align: right;">860.00</td> <td style="width:50%;"></td> </tr> <tr> <td style="text-align: right;">\$</td> <td style="text-align: right;">130.00</td> <td></td> </tr> </table>		\$	860.00		\$	130.00	
\$	860.00								
\$	130.00								
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 C.F.R. 1.492(e)).									
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE						
Total Claims	54	-20 = 34	X \$18.00						
Independent Claims	13	-3 = 10	X \$80.00						
MULTIPLE DEPENDENT CLAIMS(S) (if applicable)			\$270.00						
TOTAL OF ABOVE CALCULATIONS =			\$ 2402.00						
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.			0.00						
SUBTOTAL =			\$ 2402.00						
Processing fee of \$130.00, for furnishing the English Translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 C.F.R. 1.492(f)).			0.00						
TOTAL NATIONAL FEE =			\$ 2402.00						
Fee for recording the enclosed assignment (37 C.F.R. 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. 3.28, 3.31). \$40.00 per property			0.00						
Fee for Petition to Revive Unintentionally Abandoned Application (\$1240.00 - Small Entity = \$620.00)			0.00						
TOTAL FEES ENCLOSED =			\$ 2402.00						
			Amount to be:						
			refunded \$						
			Charged \$						

- a. ☒ A check in the amount of \$2402.00 to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. 14-1140 in the amount of \$_____ to cover the above fees. A duplicate copy of this form is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 14-1140. A duplicate copy of this form is enclosed.
- d. ☒ The entire content of the foreign application(s), referred to in this application is/are hereby incorporated by reference in this application.

NOTE: Where an appropriate time limit under 37 C.F.R. 1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

NIXON & VANDERHYE P.C.
 1100 North Glebe Road, 8th Floor
 Arlington, Virginia 22201-4714
 Telephone: (703) 816-4000



 SIGNATURE

B. J. Sadoff

 NAME


36,663 **August 10, 2001**

 REGISTRATION NUMBER Date

#7

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

SCHREIBERAtty. Ref.: **35-213**Serial No. **09/913,159**Group: **Unknown**Filed: **August 10, 2001**Examiner: **Unknown****VIRAL VACCINE**

* * * * *

March 11, 2002Assistant Commissioner for Patents
Washington, DC 20231

Sir:

AMENDMENT

Responsive to the Notification dated January 18, 2002, and further to the Request of February 6, 2002, entry and consideration of the following amendments and remarks are requested.

IN THE SPECIFICATION

Amend the specification as follows:

Insert the attached Sequence Listing in place of the copy of the same filed August 10, 2001.

REMARKS

Reconsideration is requested.

RECEIVED 03/11/02

A copy of the Notification of Missing Requirements dated January 18, 2002, and Raw Sequence Listing Error Report attached to the same, are attached. The specification has been amended to include the attached Sequence Listing.

The attached paper and computer-readable copies of the Sequence Listing are the same. No new matter has been added. A separate Statement to this effect is attached.


Nothing further is believed to be required in response to the Notification of Missing Requirements dated January 18, 2002, however the Office is requested to contact the undersigned if otherwise.

An early and favorable Action on the merits is requested.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: _____



B. J. Sadoff

Reg. No. 36,663

BJS:eaw

1100 North Glebe Road, 8th Floor
Arlington, VA 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100

#7

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

SCHREIBER

Atty. Ref.: **35-213**

Serial No. **09/913,159**

Group: **Unknown**

Filed: **August 10, 2001**

Examiner: **Unknown**



VIRAL VACCINE

* * * * *

March 11, 2002

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

STATEMENT

The attached paper and computer-readable copies of the Sequence Listing are the same. No new matter has been added.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: _____

B. J. Sadoff

B. J. Sadoff
Reg. No. **36,663**

BJS:eaw

1100 North Glebe Road, 8th Floor
Arlington, VA 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100

RECEIVED MAR 11 2002



PCT09

ENTERED

RAW SEQUENCE LISTING

DATE: 04/29/2002

PATENT APPLICATION: US/09/913,159A

TIME: 11:20:40

Input Set : A:\seq listing.txt

Output Set: N:\CRF3\04292002\I913159A.raw

4 <110> APPLICANT: Strathmann AG & Co.
 6 <120> TITLE OF INVENTION: Virus-Vaccine
 8 <130> FILE REFERENCE: P057760
 C--> 10 <140> CURRENT APPLICATION NUMBER: US/09/913,159A
 C--> 11 <141> CURRENT FILING DATE: 2002-03-11
 13 <150> PRIOR APPLICATION NUMBER: 199 07 485.2
 14 <151> PRIOR FILING DATE: 1999-02-12
 16 <160> NUMBER OF SEQ ID NOS: 12
 18 <170> SOFTWARE: PatentIn Ver. 2.1
 20 <210> SEQ ID NO: 1
 21 <211> LENGTH: 9709
 22 <212> TYPE: DNA
 23 <213> ORGANISM: Human immunodeficiency virus
 25 <400> SEQUENCE: 1
 26 tggaaagggct aatttggtcc caaaaaagac aagagatcct tgatctgtgg atctaccaca 60
 27 cacaaggcta ctccctgat tggcagaact acacaccagg gccagggatc agatatccac 120
 28 tgaccttttg atggtgcttc aagtttagtac cagttgaacc agagcaagta gaagaggcca 180
 29 aataaggaga gaagaacagc ttgttacacc ctatgagcca gcatgggatg gaggaccgg 240
 30 agggagaagt attagtgtgg aagtttgaca gcctcctagc atttcgtcac atggcccag 300
 31 agctgcatcc ggagtactac aaagactgct gacatcgagc tttctacaag ggactttccg 360
 32 ctggggactt tccagggagg tgtggcctgg gcgggactgg ggagtggcga gccctcagat 420
 33 gctacatata agcagctgct ttttgctgt actgggtctc tctggttaga ccagatctga 480
 34 gctggggagc tctctggcta actaggaac ccaactgctta agcctcaata aagcttgcc 540
 35 tgagtgtctca aagtagtgtg tgcccgtctg ttgtgtgact ctggtaacta gagatccctc 600
 36 agaccctttt agtcagttg gaaaatctct agcagtggcg cccgaacagg gacttgaag 660
 37 cgaaagtaaa gccagaggag atctctcgac gcaggactcg gcttgctgaa gcgcgcacgg 720
 38 caagaggcga gggcgggcga ctggtgagta gcgcaaaaat tttgactagc ggaggctaga 780
 39 aggagagaga tgggtgcgag agcgtcggtt ttaagcgggg gagaattaga taaatgggaa 840
 40 aaaattcggg taaggccagg gggaaagaaa caatataaac taaaacatat agtatgggca 900
 41 agcagggagc tagaacgatt cgcagttaat cctggccttt tagagacatc agaaggctgt 960
 42 agacaaatac tgggacagct acaaccatcc cttcagacag gatcagaaga acttagatca 1020
 43 ttatataata caatagcagt cctctattgt gtgcatcaaa ggatagatgt aaaagacacc 1080
 44 aaggaagcct tagataagat agaggaagag caaaacaaaa gtaagaaaaa ggcacagcaa 1140
 45 gcagcagctg acacaggaaa caacagccag gtcagccaaa attaccctat agtcagaaac 1200
 46 ctccaggggc aaatggtaca tcaggccata tcacctagaa ctttaaatgc atgggtaaaa 1260
 47 gtagtagaag agaaggcttt cagcccagaa gtaataccca tgttttcagc attatcagaa 1320
 48 ggagccaccc cacaagattt aaataccatg ctaaaccacag tggggggaca tcaagcagcc 1380
 49 atgcaaatgt taaaagagac catcaatgag gaagctgcag aatgggatag attgcatcca 1440
 50 gtgcatgcag ggctatttgc accaggccag atgagagaac caaggggaag tgacatagca 1500
 51 ggaactacta gtacccttca ggaacaaata ggaatggaac cacataatcc acctatccca 1560
 52 gtaggagaaa tctataaaag atggataatc ctgggattaa ataaaaatag aagaatgtat 1620
 53 agccctacca gcattctgga cataagacaa ggaccaaagg aaccctttag agactatgta 1680
 54 gaccgattct ataaaaactc aagagccgag caagcttcac aagaggtaaa aaattggatg 1740

RAW SEQUENCE LISTING

DATE: 04/29/2002

PATENT APPLICATION: US/09/913,159A

TIME: 11:20:40

Input Set : A:\seq_listing.txt

Output Set: N:\CRF3\04292002\I913159A.raw

T00T60"CAT660

```

55 acagaaacct tgttgggtcca aaatgcgaac ccagattgta agactatttt aaaagcattg 1800
56 ggaccaggag cgacactaga agaaatgatg acagcatgtc agggagtggg gggacccggc 1860
57 cataaagcaa gagttttggc tgaagcaatg agccaagtaa caaatccagc taccataatg 1920
58 atacagaaaag gcaattttag gaaccaaaga aagactgtta agtgtttcaa ttgtggcaaa 1980
59 gaaggggcaca tagccaaaaa ttgcagggcc cctaggaaaa agggctgttg gaaatgtgga 2040
60 aaggaaggac accaaatgaa agattgtact gagagacagg ctaatttttt agggaagatc 2100
61 tggccttccc acaagggaag gccagggaat tttcttcaga gcagaccaga gccaacagcc 2160
62 ccaccagaag agagcttcag gtttggggaa gagacaacaa ctccctctca gaagcaggag 2220
63 ccgatagaca aggaactgta tccttttagct tccctcagat cactcttttg cagcgacccc 2280
64 tcgtcacaat aaagataggg gggcaattaa aggaagctct attagataga ggagcagatg 2340
65 atacagtatt agaagaaatg aatttgcag gaagatggaa accaaaaatg atagggggaa 2400
66 ttggagggtt tatcaaagta ggacagtatg atcagatact catagaaatc tgcggacata 2460
67 aagctatagg tacagtatta gtaggaccta cacctgtcaa cataattgga agaaatctgt 2520
68 tgactcagat tggctgcact ttaaattttc cctagtctcc tattgagact gtaccagtaa 2580
69 aattaaagcc aggaatggat ggccaaaag ttaaacaatg gccattgaca gaagaaaaaa 2640
70 taaaagcatt agtagaatt tgtacagaaa tggaaaagga aggaaaaatt tcaaaaattg 2700
71 ggctgaaaa tccatacaat actccagtat ttgccataaa gaaaaaagac agtactaaat 2760
72 ggagaaaaat agtagatttc agagaactta ataagagaa tcaagatttc tgggaagttc 2820
73 aattaggaat accacatcct gcagggttaa aacagaaaaa atcagtaaca gtactggatg 2880
74 tgggcgatgc atatttttca gttcccttag ataaagactt cagggaagtat actgcattta 2940
75 ccatacctag tataaacaat gagacaccag ggattagata tcagtacaat gtgcttcac 3000
76 agggatggaa aggatcacca gcaatattcc agtgtagcat gacaaaaatc ttagagcctt 3060
77 ttagaaaaa aaatccagac atagtcatct atcaatacat ggatgatttg tatgtaggat 3120
78 ctgacttaga aatagggcag catagaacaa aaatagagga actgagacaa catctgttga 3180
79 ggtggggatt taccacacca gacaaaaaac atcagaaaga acctccattc ctttggtagg 3240
80 gttatgaact ccatcctgat aaatggacag tacagcctat agtgctgcca gaaaaggaca 3300
81 gctggactgt caatgacata cagaaattag tgggaaaaat gaattgggca agtcagattt 3360
82 atgcagggat taaagtaagg caattatgta aacttcttag gggaaccaa gcactaacag 3420
83 aagtagtacc actaacagaa gaagcagagc tagaactggc agaaaacagg gagattctaa 3480
84 aagaaccggt acatggagtg tattatgacc catcaaaaga cttaatagca gaaatacaga 3540
85 agcaggggca aggccaatgg acatatcaaa tttatcaaga gccattttaa aatctgaaaa 3600
86 caggaaaaa tgcaagaatg aagggtgccc acactaatga tgtgaaacaa ttaacagagg 3660
87 cagtacaaaa aatagccaca gaaagcatag taatatgggg aaagactcct aaattttaat 3720
88 taccataca aaaggaacaa tgggaagcat ggtggacaga gtattggcaa gccacctgga 3780
89 ttctgagtg ggagtttgtc aataccctc ccttagtgaa gttatggtac cagttagaga 3840
90 aagaacccat aataggagca gaaactttct atgtagatgg ggcagccaat agggaaacta 3900
91 aattaggaag agcaggatat gtaactgaca gaggaagaca aaaagttgtc cccctaacgg 3960
92 acacaacaaa tcagaagact gagttacaag caattcatct agctttgcag gattcgggat 4020
93 tagaagtaaa catagtgaac gactcacaat atgcattggg aatcattcaa gcacaaccag 4080
94 ataagagtga atcagagtta gtcagtcaaa taatagagca gtttaataaa aaggaaaaag 4140
95 tctacctggc atgggtacca gcacacaaag gaattggagg aaatgaacaa gtagatgggt 4200
96 tggtcagtgc tggaaatcagg aaagtactat ttttagatgg aatagataag gcccaagaag 4260
97 aacatgagaa atatcacagt aattggagag caatggctag tgattttaac ctaccacctg 4320
98 tagtagcaaa agaaatagta gccagctgtg ataaatgtca gctaaaaggg gaagccatgc 4380
99 atggacaagt agactgtac ccaggaatat ggcagctaga ttgtacacat ttagaaggaa 4440
100 aagttatctt gtagcagtt catgtagcca gtggatatat agaagcagaa gtaattccag 4500
101 cagagacagg gcaagaaaca gcatacttcc tcttaaaatt agcaggaaga tggccagtaa 4560
102 aaacagtaca tacagacaat ggcagcaatt tcaccagtac tacagttaag gccgctgtt 4620
103 ggtggggcgg gatcaagcag gaatttggca tccctacaa tcccaaagt caaggagtaa 4680

```

RAW SEQUENCE LISTING

DATE: 04/29/2002

PATENT APPLICATION: US/09/913,159A

TIME: 11:20:40

Input Set : A:\seq listing.txt

Output Set: N:\CRF3\04292002\I913159A.raw

104 tagaatctat gaataaagaa ttaaagaaaa ttataggaca ggtaagagat caggctgaac 4740
105 atcttaagac agcagtacaa atggcagtat tcatccacaa ttttaaaaga aaagggggga 4800
106 ttgggggggta cagtgcaggg gaaagaatag tagacataat agcaacagac atacaaacta 4860
107 aagaattaca aaaacaaatt acaaaaattc aaaattttcg ggtttattac agggacagca 4920
108 gagatccagt ttggaaagga ccagcaaagc tcctctggaa aggtgaaggg gcagtagtaa 4980
109 tacaagataa tagtgacata aaagtagtgc caagaagaaa agcaaagatc atcagggatt 5040
110 atggaaaaca gatggcaggt gatgattgtg tggcaagtag acaggatgag gattaacaca 5100
111 tggaaaagat tagtaaaaca ccatatgtat atttcaagga aagctaagga ctggttttat 5160
112 agacatcact atgaaagtac taatccaaaa ataagttcag aagtacacat cccactaggg 5220
113 gatgctaaat tagtaataac aacatattgg ggtctgcata caggagaaaag agactggcat 5280
114 ttgggtcagg gagtctccat agaattggag aaaaagagat atagcacaca agtagaccct 5340
115 gacctagcag accaactaat tcatctgcac tatttttgatt gtttttcaga atctgctata 5400
116 agaaatacca tcttaggacg tatagttagt cctaggtgtg aatatcaagc aggcataaac 5460
117 aaggtaggat ctctacagta cttggcacta gcagcattaa taaaaccaa acagataaag 5520
118 ccacctttgc ctagtgttag gaaactgaca gaggacagat ggaacaagcc ccagaagacc 5580
119 aagggccaca gagggagcca tacaatgaat ggacactaga gcttttagag gaacttaaga 5640
120 gtgaagctgt tagacatttt ctaggatata ggctccataa cttaggacaa catatctatg 5700
121 aaacttacgg ggatacttgg gcaggagtgg aagccataat aagaattctg caacaactgc 5760
122 tgtttatcca tttcagaatt ggggtctgac atagcagaat aggcgttact cgacagagga 5820
123 gagcaagaaa tggagccagt agatccctaga ctagagccct ggaagcatcc aggaagtcag 5880
124 cctaaaactg cttgtaccaaa ttgctattgt aaaaagtgtt gctttcattg ccaagtttgt 5940
125 ttcattgaca aagccttagg catctcctat ggcaggaaga agcggagaca gcgacgaaga 6000
126 gctcatcaga acagtcagac tcatcaagct tctctatcaa agcagtaagt agtacatgta 6060
127 atgcaaccta taatagtagc aatagtagca ttagtagtag caataataat agcaatagtt 6120
128 gtgtggtcca tagtaatcat agaatatagg aaaatattaa gacaaagaaa aatagacagg 6180
129 ttaattgata gactaataga aagagcagaa gacagtggca atgagagtga aggagaagta 6240
130 tcagcacttg tggagatggg ggtggaaaatg gggcaccatg ctcccttggga tattgatgat 6300
131 ctgtagtgct acagaaaaat tgtgggtcac agtctattat ggggtacctg tgtggaagga 6360
132 agcaaccacc actctatttt gtgcatcaga tgctaaagca tatgatacag aggtacataa 6420
133 tgttttgggc acacatgcct gtgtacccac agaccccaac ccacaagaag tagtattggt 6480
134 aaatgtgaca gaaaatttta acatgtggaa aatgacatg gtagaacaga tgcattgagga 6540
135 tataatcagt ttatgggata aaagcctaaa gccatgtgta aaattaacc cactctgtgt 6600
136 tagtttaag tgcactgatt tgaagaatga tactaatacc aatagtagta gcgggagaat 6660
137 gataatggag aaaggagaga taaaaaactg ctctttcaat atcagcacaa gcataagaga 6720
138 taagggtcag aaagaatatg cattctttta taaacttgat atagtaccaa tagataatac 6780
139 cagctatagg ttgataagtt gtaaacacct agtcattaca caggcctgtc caaaggatc 6840
140 ctttgagcca attcccatac attattgtgc cccggctggg tttgcgattc taaaatgtaa 6900
141 taataagacg ttcaatggaa caggaccatg taaaaatgtc agcacagtac aatgtacaca 6960
142 tggaaatcagg ccagtagtat caactcaact gctgtttaat ggcagtctag cagaagaaga 7020
143 tgtagtaatt agatctgcca atttcacaga caatgctaaa accataatag tacagctgaa 7080
144 cacatctgta gaaattaatt gtacaagacc caacaacaat acaagaaaaa gtatccgtat 7140
145 ccagagggga ccagggagag catttggttac aataggaaaa ataggaaata tgagacaagc 7200
146 acattgtaac attagtagag caaaatggaa tgccacttta aaacagatag ctagcaaat 7260
147 aagagaacaa tttggaaata ataaaacaat aatctttaag caatcctcag gaggggaccc 7320
148 agaaattgta acgcacagtt ttaattgtgg aggggaattt ttctactgta attcaacaca 7380
149 actgtttaat agtacttggt ttaatagtac ttggagtact gaagggtcaa ataactga 7440
150 aggaagtgc acaatcacac tcccatgcag aataaaacaa tttataaaca tgtggcagga 7500
151 agtaggaaaa gcaatgtatg cccctcccat cagtggacaa attagatgtt catcaaatat 7560
152 tactgggctg ctattaacaa gagatggtgg taataacaac aatgggtccg agatcttcag 7620

RAW SEQUENCE LISTING

DATE: 04/29/2002

PATENT APPLICATION: US/09/913,159A

TIME: 11:20:40

Input Set : A:\seq listing.txt

Output Set: N:\CRF3\04292002\I913159A.raw

```

153 acctggagga ggcgatatga gggacaattg gagaagtga ttatataaat ataaagtagt 7680
154 aaaaattgaa ccattaggag tagcaccac caaggcaaag agaagagtgg tgcagagaga 7740
155 aaaaagagca gtgggaatag gagctttgtt ccttgggttc ttgggagcag caggaagcac 7800
156 tatgggctgc acgtcaatga cgctgacggt acaggccaga caattattgt ctgatatagt 7860
157 gcagcagcag aacaatttgc tgagggctat tgaggcgcaa cagcatctgt tgcaactcac 7920
158 agtctggggc atcaaacagc tccaggcaag aatcctggct gtggaaagat acctaaagga 7980
159 tcaacagctc ctggggattt ggggttgctc tggaaaactc atttgacca ctgctgtgcc 8040
160 ttggaatgct agttggagta ataaatctct ggaacagatt tggaataaca tgacctggat 8100
161 ggagtgggac agagaaatta acaattacac aagcttaata cactccttaa ttgaagaatc 8160
162 gcaaaaccag caagaaaaga atgaacaaga attattggaa ttagataaat gggcaagttt 8220
163 gtggaattgg tttaacataa caaattggct gtggtatata aaattattca taatgatagt 8280
164 aggaggcttg gtaggtttta gaatagtttt tgctgtactt tctatagtga atagagttag 8340
165 gcagggatat tcaccattat cgtttcagac ccacctccca atcccagggg gaccgcagac 8400
166 gcccgaagga atagaagaag aaggtggaga gagagacaga gacagatcca ttcgattagt 8460
167 gaacggatcc ttagctacta tctgggacga tctgcgagc ctgtgcctct tcagctacca 8520
168 ccgcttgaga gacttactct tgattgtaac gagattgtg gaacttctg gacgcagggg 8580
169 gtgggaagcc ctcaaattat ggtggaatct cctacagtat tggagtcagg aactaaagaa 8640
170 tagtgctgtt aacttgctca atgccacagc catagcagta gctgagggga cagatagggt 8700
171 tatagaagta ttacaagcag cttatagagc tattgocac atacctagaa gaataagaca 8760
172 gggcttggaaggattttgc tataagatgg gtggcaagtgt gtcaaaaagt agtgtgattg 8820
173 gatggcctgc tgtaaggga agaagtagac gagctgagcc agcagcagat ggggtgggag 8880
174 cagtatctcg agacctagaa aaacatggag caatcacaag tagcaatata gcagctaaca 8940
175 atgctgcttg tgcctggcta gaagcacaag aggaggaaga ggtgggtttt ccagtcacac 9000
176 ctgaggtacc ttttaagacca atgacttaca aggcagctgt agatcttagc cactttttta 9060
177 aagaaaaggg gggactggaa gggctaattc actcccaaag aagacaagat atccttgatc 9120
178 tgtggatcta ccacacacaa ggctacttcc ctgattggca gaactacaca ccagggccag 9180
179 gggtcagata tccactgacc tttggatggt gctacaagct agtaccagtt gagccagata 9240
180 aggtagaaga ggccaataaa ggagagaaca ccagottggt acaccctgtg agcctgcatg 9300
181 gaatggatga ccctgagaga gaagtgttag agtggagggt tgacagccgc ctagcatttc 9360
182 atcacgtggc ccgagagctg catccggagt acttcaagaa ctgctgacat cgagcttgct 9420
183 acaagggact ttccgctggg gactttccag ggaggcgtgg cctgggcggg actggggagt 9480
184 ggcgagccct cagatgctgc atataagcag ctgctttttg cctgtactgg gtctctctgg 9540
185 ttagaccaga tctgagcctg ggagctctct ggctaactag ggaaccact gcttaagcct 9600
186 caataaagct tgcttgagt gcttcaagta gtgtgtgcc gtctgttggt tgactctggt 9660
187 aactagagat ccctcagacc ctttttagtca gtgtggaaaa tctctagca 9709

```

191 <210> SEQ ID NO: 2

192 <211> LENGTH: 854

193 <212> TYPE: PRT

194 <213> ORGANISM: Human immunodeficiency virus

196 <220> FEATURE:

197 <223> OTHER INFORMATION: Envelope Polyprotein

199 <400> SEQUENCE: 2

```

200 Met Arg Val Lys Glu Lys Tyr Gln His Leu Trp Arg Trp Gly Trp Lys
201   1           5           10           15
203 Trp Gly Thr Met Leu Leu Gly Ile Leu Met Ile Cys Ser Ala Thr Glu
204           20           25           30
206 Lys Leu Trp Val Thr Val Tyr Tyr Gly Val Pro Val Trp Lys Glu Ala
207           35           40           45
209 Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys Ala Tyr Asp Thr Glu

```

DATE: 04/29/2002

TIME: 11:20:40

Output Set: N:\CRF3\04292002\I913159A.raw

210	50					55					60							
212	Val	His	Asn	Val	Trp	Ala	Thr	His	Ala	Cys	Val	Pro	Thr	Asp	Pro	Asn		
213	65						70						75	80				
215	Pro	Gln	Glu	Val	Val	Leu	Val	Asn	Val	Thr	Glu	Asn	Phe	Asn	Met	Trp		
216						85						90	95					
218	Lys	Asn	Asp	Met	Val	Glu	Gln	Met	His	Glu	Asp	Ile	Ile	Ser	Leu	Trp		
219						100						105	110					
221	Asp	Gln	Ser	Leu	Lys	Pro	Cys	Val	Lys	Leu	Thr	Pro	Leu	Cys	Val	Ser		
222						115						120	125					
224	Leu	Lys	Cys	Thr	Asp	Leu	Lys	Asn	Asp	Thr	Asn	Thr	Asn	Ser	Ser	Ser		
225						130						135	140					
227	Gly	Arg	Met	Ile	Met	Glu	Lys	Gly	Glu	Ile	Lys	Asn	Cys	Ser	Phe	Asn		
228	145						150						155	160				
230	Ile	Ser	Thr	Ser	Ile	Arg	Asp	Lys	Val	Gln	Lys	Glu	Tyr	Ala	Phe	Phe		
231						165						170	175					
233	Tyr	Lys	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Thr	Ser	Tyr	Arg	Leu	Ile		
234						180						185	190					
236	Ser	Cys	Asn	Thr	Ser	Val	Ile	Thr	Gln	Ala	Cys	Pro	Lys	Val	Ser	Phe		
237						195						200	205					
239	Glu	Pro	Ile	Pro	Ile	His	Tyr	Cys	Ala	Pro	Ala	Gly	Phe	Ala	Ile	Leu		
240						210						215	220					
242	Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys	Thr	Asn	Val		
243	225						230						235	240				
245	Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val	Ser	Thr	Gln		
246						245						250	255					
248	Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Asp	Val	Val	Ile	Arg	Ser		
249						260						265	270					
251	Ala	Asn	Phe	Thr	Asp	Asn	Ala	Lys	Thr	Ile	Ile	Val	Gln	Leu	Asn	Thr		
252						275						280	285					
255	Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr	Arg	Lys	Ser		
256						290						295	300					
258	Ile	Arg	Ile	Gln	Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Thr	Ile	Gly	Lys		
259	305						310						315	320				
261	Ile	Gly	Asn	Met	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Ala	Lys	Trp		
262						325						330	335					
264	Asn	Ala	Thr	Leu	Lys	Gln	Ile	Ala	Ser	Lys	Leu	Arg	Glu	Gln	Phe	Gly		
265						340						345	350					
267	Asn	Asn	Lys	Thr	Ile	Ile	Phe	Lys	Gln	Ser	Ser	Gly	Gly	Asp	Pro	Glu		
268						355						360	365					
270	Ile	Val	Thr	His	Ser	Phe	Asn	Cys	Gly	Gly	Glu	Phe	Phe	Tyr	Cys	Asn		
271						370						375	380					
273	Ser	Thr	Gln	Leu	Phe	Asn	Ser	Thr	Trp	Phe	Asn	Ser	Thr	Trp	Ser	Thr		
274	385						390						395	400				
276	Glu	Gly	Ser	Asn	Asn	Thr	Glu	Gly	Ser	Asp	Thr	Ile	Thr	Leu	Pro	Cys		
277						405						410	415					
279	Arg	Ile	Lys	Gln	Phe	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met		
280						420						425	430					
282	Tyr	Ala	Pro	Pro	Ile	Ser	Gly	Gln	Ile	Arg	Cys	Ser	Ser	Asn	Ile	Thr		
283						435						440	445					

RAW SEQUENCE LISTING ERROR SUMMARY
PATENT APPLICATION: US/09/913,159A

DATE: 04/29/2002
TIME: 11:20:41

Input Set : A:\seq listing.txt
Output Set: N:\CRF3\04292002\I913159A.raw

Please Note:

Use of n and/or Xaa have been detected in the Sequence Listing. Please review the Sequence Listing to ensure that a corresponding explanation is presented in the <220> to <223> fields of each sequence which presents at least one n or Xaa.

Seq#:4; N Pos. 99

FOOTNOTES CONTAINED

VERIFICATION SUMMARY

DATE: 04/29/2002

PATENT APPLICATION: US/09/913,159A


TIME: 11:20:41

Input Set : A:\seq listing.txt

Output Set: N:\CRF3\04292002\I913159A.raw

L:10 M:270 C: Current Application Number differs, Replaced Application Number
L:11 M:271 C: Current Filing Date differs, Replaced Current Filing Date
L:399 M:341 W: (46) "n" or "Xaa" used, for SEQ ID#:4 after pos.:60

T00T60"6aT60



11 MAR 2002

#7

SEQUENCE LISTING

<110> Strathmann AG & Co.

<120> Virus-Vaccine

<130> P057760

<140>

<141>

<150> 199 07 485.2

<151> 1999-02-12

<160> 12

<170> PatentIn Ver. 2.1

<210> 1

<211> 9709

<212> DNA

<213> Human immunodeficiency virus

<400> 1

tggaagggct aatttggtcc caaaaaagac aagagatcct tgatctgtgg atctaccaca 60
cacaaggcta cttccctgat tggcagaact acacaccagg gccagggatc agatatccac 120
tgaccttttg atggtgcttc aagtttagtac cagttgaacc agagcaagta gaagaggcca 180
aataaggaga gaagaacagc ttgttacacc ctatgagcca gcatgggatg gaggaccccg 240
agggagaagt attagtgtgg aagtttgaca gcctcctagc atttcgtcac atggcccgag 300
agctgcatcc ggagtactac aaagactgct gacatcgagc tttctacaag ggactttccg 360
ctggggactt tccagggagg tgtggcctgg gcgggactgg ggagtggcga gccctcagat 420
gctacatata agcagctgct ttttgccctgt actgggtctc tctgggttaga ccagatctga 480
gcctggggagc tctctggcta actaggggaac cactgctta agcctcaata aagcttgcct 540
tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact ctggtaacta gagatccctc 600
agaccctttt agtcagtggt gaaaatctct agcagtggcg cccgaacagg gacttgaaag 660
cgaaagtaaa gccagaggag atctctcgac gcaggactcg gcttgttgaa gcgcgcacgg 720
caagaggcga ggggcggcga ctggtgagta cgccaaaaat tttgactagc ggaggctaga 780
aggagagaga tgggtgagag agcgtcggtt ttaagcgggg gagaattaga taaatgggaa 840
aaaattcggg taaggccagg gggaaagaaa caatataaac taaaacatat agtatgggca 900
agcaggggagc tagaacgatt cgcagttaat cctggccttt tagagacatc agaaggctgt 960
agacaaatac tgggacagct acaaccatcc ctccagacag gatcagaaga acttagatca 1020
ttatataata caatagcagt cctctattgt gtgcatcaaa ggatagatgt aaaagacacc 1080
aaggaagcct tagataagat agaggaagag caaaacaaaa gtaagaaaaa ggcacagcaa 1140
gcagcagctg acacaggaaa caacagccag gtcagccaaa attaccctat agtgcagaac 1200
ctccaggggc aaatggtaca tcaggccata tcacctagaa ctttaaattgc atgggtaaaa 1260
gtagtagaag agaaggcttt cagcccagaa gtaataccca tgttttcagc attatcagaa 1320
ggagccaccc cacaagattt aaataccatg ctaaacacag tgggggggaca tcaagcagcc 1380
atgcaaagt taaaagagac catcaatgag gaagctgcag aatgggatag attgcatcca 1440
gtgcatgcag ggcctattgc accaggccag atgagagAAC caaggggaag tgacatagca 1500
ggaactacta gtacccttca ggaacaaata ggatggatga cacataatcc acctatccca 1560
gtaggagaaa tctataaaag atggataatc ctgggattaa ataaaatagt aagaatgtat 1620
agccctacca gcattctgga cataagacaa ggaccaaagg aaccctttag agactatgta 1680
gaccgattct ataaaactct aagagccgag caagcttcac aagaggtaaa aaattggatg 1740
acagaaacct tgttgggtcca aaatgcgaac ccagattgta agactatttt aaaagcattg 1800
ggaccaggag cgacactaga agaaatgatg acagcatgtc agggagtggg gggacccggc 1860
cataaagcaa gagttttggc tgaagcaatg agccaagtaa caaatccagc taccataatg 1920

100760-57760

atacagaaag	gcaatttttag	gaaccaaaaga	aagactgttta	agtgtttcaa	ttgtggcaaa	1980
gaagggcaca	tagccaaaaa	ttgcagggcc	cctaggaaaa	agggctgttg	gaaatgtgga	2040
aaggaaggac	accaaataaa	agattgtact	gagagacagg	ctaatttttt	aggggaagatc	2100
tggccttccc	acaaggggaag	gccaggggaat	tttcttcaga	gcagaccaga	gccaacagcc	2160
ccaccagaag	agagcttcag	gtttggggaa	gagacaacaa	ctccctctca	gaagcaggag	2220
ccgatagaca	aggaactgta	tccttttagct	tccttcagat	cactcttttg	cagcgacccc	2280
tcgtcacaa	aaagataggg	gggcaattaa	aggaagctct	attagataca	ggagcagatg	2340
atacagtatt	agaagaaatg	aatttggccag	gaagatggaa	acaaaaaatg	ataggggggaa	2400
ttggagggtt	tatcaaagta	ggacagtatg	atcagatact	catagaaatc	tgcggacata	2460
aagctatagg	tacagtatta	gtaggaccta	cacctgtcaa	cataattgga	agaaatctgt	2520
tgactcagat	tggctgcact	ttaaattttc	ccattagtcc	tattgagact	gtaccagtaa	2580
aattaaagcc	aggaatggat	ggccaaaaag	ttaaacaatg	gccattgaca	gaagaaaaaa	2640
taaaagcatt	agtagaaatt	tgtacagaaa	tggaaaagga	aggaaaaatt	tcaaaaattg	2700
ggcctgaaaa	tccatacaat	actccagtat	ttgccataaa	gaaaaaagac	agtactaaat	2760
ggagaaaatt	agtagatttc	agagaactta	ataagagaa	tcaagatttc	tgggaagtgc	2820
aattaggaat	accacatcct	gcagggttaa	aacagaaaaa	atcagtaaca	gtactggatg	2880
tgggcgatgc	atatttttca	gttcccttag	ataaagactt	caggaagtat	actgcattta	2940
ccatacctag	tataaacaat	gagacaccag	ggattagata	tcagtacaat	gtgcttccac	3000
agggatggaa	aggtaccaca	gcaatatctc	agtgtagcat	gacaaaaatc	ttagagcctt	3060
ttagaaaaaa	aaatccgagc	atagtcatct	atcaatacat	ggatgatttg	tatgtaggat	3120
ctgacttaga	aatagggcag	catagaacaa	aaatagagga	actgagacaa	catctgttga	3180
ggtggggatt	taccacacca	gacaaaaaac	atcagaaaga	acctccattc	ctttgggatg	3240
gttatgaact	ccatcctgat	aaatggacag	tacagcctat	agtgtctgca	gaaaaggaca	3300
gctggactgt	caatgacata	cagaaattag	tgggaaaatt	gaattgggca	agtcagattt	3360
atgcagggat	taaagtaagg	caattatgta	aacttcttag	gggaaccaa	gcactaacag	3420
aagtagtacc	actaacagaa	gaagcagagc	tagaactggc	agaaaacagg	gagattctaa	3480
agaaccgggt	acatggagtg	tattatgacc	catcaaaaga	cttaatagca	gaaatacaga	3540
agcaggggca	aggccaatgg	acatatcaaa	tttatcaaga	gccattttaa	aatctgaaaa	3600
caggaaaata	tgcaagaatg	aagggtgccc	acactaatga	tgtgaaacaa	ttaacagagg	3660
cagtacaaaa	aatagccaca	gaaagcatag	taatatgggg	aaagactcct	aaattttaat	3720
taccataaca	aaaggaaaca	tgggaagcat	ggtggacaga	gtattggcaa	gccacctgga	3780
ttcttgagtg	ggagtttgtc	aatacccctc	ccttagtgaa	gttatggtac	cagttagaga	3840
aagaacccat	aataggagca	gaaactttct	atgtagatgg	ggcagccaat	agggaaacta	3900
aattagggaa	agcaggatat	gtaactgaca	gaggaagaca	aaaagttgtc	cccctaacgg	3960
acacaacaaa	tcagaagact	gagttacaag	caattcatct	agctttgcag	gattcgggat	4020
tagaagtaaa	catagtgcac	gactcacaat	atgcattggg	aatcattcaa	gcacaaccag	4080
ataagatgta	atcagagtta	gtcagtcaca	taatagagca	gttaataaaa	aaggaaaaag	4140
tctacctggc	atgggtacca	gcacacaaag	gaattggagg	aaatgaacaa	gtagatgggt	4200
tggtcagtgc	tggaaatcagg	aaagtactat	ttttagatgg	aatatgataag	gcccaagaag	4260
aacatgagaa	atatcacagt	aattggagag	caatggctag	tgatttttaac	ctaccacctg	4320
tagtagcaaa	agaaatagta	gccagctgtg	ataaatgtca	gctaaaaggg	gaagccatgc	4380
atggacaagt	agactgtagc	ccaggaatat	ggcagctaga	ttgtacacat	ttagaaggaa	4440
aagtattctt	ggtagcagtt	catgtagcca	gtggatatat	agaagcagaa	gtaattccag	4500
cagagacagg	gcaagaaaca	gcatacttcc	tcttaaaatt	agcaggaaga	tggccagtaa	4560
aaacagtaca	tacagacaat	ggcagcaatt	tcaccagtac	tacagttaag	gccgcctggt	4620
ggtgggcggg	gatcaagcag	gaatttggca	ttccctacaa	tcccaaagt	caaggagtaa	4680
tagaatctat	gaataaagaa	ttaaagaaaa	ttataggaca	ggtaagagat	caggctgaac	4740
atcttaagac	agcagtacaa	atggcagtat	tcattccaaa	ttttaaaaga	aaagggggga	4800
ttggggggta	cagtgcaggg	gaaagaatag	tagacataat	agcaacagac	atacaaaacta	4860
aagaattaca	aaaacaaatt	acaaaaattc	aaaattttcg	ggtttattac	agggacagca	4920
gagatccagt	ttggaaaagga	ccagcaaagc	tcctctggaa	aggtgaaggg	gcagtagtaa	4980
tacaagataa	tagtgacata	aaagtagtgc	caagaagaaa	agcaaagatc	atcagggatt	5040
atggaaaaca	gatggcaggt	gatgatgtgt	tggcaagtag	acaggatgag	gattaacaca	5100
tggaaaagat	tagtaaaaaa	ccatatgtat	atttcaagga	aagctaagga	ctggttttat	5160
agacatcact	atgaaagtac	taatccaaaa	ataagttcag	aagtacacat	cccactaggg	5220
gatgctaaat	tagtaataac	aacatattgg	ggtctgcata	caggagaaag	agactggcat	5280
ttgggtcagg	gagtctccat	agaatggagg	aaaaagagat	atagcacaca	agtagaccct	5340

gacctagcag	accaactaat	tcatctgcac	tatttttgatt	gtttttcaga	atctgctata	5400
agaaatacca	tattaggacg	tatagttagt	cctagggtgtg	aatatcaagc	aggacataac	5460
aaggtaggat	ctctacagta	cttggcacta	gcagcattaa	taaaaccaa	acagataaag	5520
ccacctttgc	ctagtgttag	gaaactgaca	gaggacagat	ggaacaagcc	ccagaagacc	5580
aagggccaca	gagggagcca	tacaatgaat	ggacactaga	gcttttagag	gaacttaaga	5640
gtgaagctgt	tagacatttt	cctaggatat	ggctccataa	cttaggacaa	catatctatg	5700
aaacttacgg	ggatacttgg	gcaggagtgg	aagccataat	aagaattctg	caacaactgc	5760
tgtttatcca	tttcagaatt	gggtgtcgac	atagcagaat	aggcgttact	cgacagagga	5820
gagcaagaaa	tggagccagt	agatcctaga	ctagagccct	ggaagcatcc	aggaagtcag	5880
cctaaaactg	ctgtaccaaa	ttgctattgt	aaaagtgtt	gctttcattg	ccaagtttgt	5940
ttcatgacaa	aagccttagg	catctcctat	ggcaggaaga	agcggagaca	gcgacgaaga	6000
gctcatcaga	acagtcagac	tcatcaagct	tctctatcaa	agcagtaagt	agtacatgta	6060
atgcaaccta	taatagtagc	aatagtagca	ttagtagtag	caataataat	agcaatagtt	6120
gtgtggtcca	tagtaatcat	agaatatagg	aaaatattaa	gacaaagaaa	aatagacagg	6180
ttaattgata	gactaataga	aagagcagaa	gacagtggca	atgagagtga	aggagaagta	6240
tcagcacttg	tggagatggg	ggtggaaatg	gggcaccatg	ctccttggga	tattgatgat	6300
ctgtagtgtc	acagaaaaat	tgtgggtcac	agtctattat	ggggtacctg	tgtggaagga	6360
agcaaccacc	actctatttt	gtgcatcaga	tgctaaagca	tatgatacag	aggtaacata	6420
tgtttgggccc	acacatgcct	gtgtacccac	agaccccaac	ccacaagaag	tagtatttgtt	6480
aaatgtgaca	gaaaatttta	acatgtggaa	aaatgacatg	gtagaacaga	tgcattgagga	6540
tataatcagt	ttatgggatc	aaagcctaaa	gcatgtgtga	aaattaaccc	cactctgtgt	6600
tagtttaaaag	tgcactgatt	tgaagaatga	tactaatacc	aatagtagta	gcggggagaat	6660
gataatggag	aaaggagaga	taaaaaactg	ctctttcaat	atcagcacia	gcataagaga	6720
taaggtgcag	aaagaatatg	cattctttta	taaacttgat	atagtaccaa	tagataatac	6780
cagctatagg	ttgataagtt	gtaacacctc	agtcattaca	caggcctgtc	caaaggtatc	6840
ctttgagcca	attcccatac	attattgtgc	ccgggtggtt	tttgcgattc	taaaatgtaa	6900
taataagacg	ttcaatggaa	caggaccatg	tacaaatgtc	agcacagtac	aatgtacaca	6960
tggaatcagg	ccagtagtat	caactcaact	gctgttaaat	ggcagtcctag	cagaagaaga	7020
tgtagttaatt	agatctgcca	atttcacaga	caatgctaaa	accataatag	tacagctgaa	7080
cacatctgta	gaaattaatt	gtacaagacc	caacaacaat	acaagaaaaa	gtatccgtat	7140
ccagagggga	ccaggggagag	catttggttac	aataggaaaa	ataggaaata	tgagacaagc	7200
acattgtaac	attagtagag	caaaatggaa	tgccacttta	aaacagatag	ctagcaaatt	7260
aagagaacaa	tttggaataa	ataaaacaat	aatctttaag	caatcctcag	gaggggaccc	7320
agaaattgta	acgcacagtt	ttaattgtgg	aggggaattt	ttctactgta	attcaacaca	7380
actgtttaat	agtacttggt	ttaatagtac	ttggagtact	gaagggtcaa	ataacactga	7440
aggaagtgc	acaatcacac	tcccatgcag	aataaaacaa	tttataaaca	tgtggcgagga	7500
agtaggaaaa	gcaatgtatg	cccctcccat	cagtggacaa	attagatggt	catcaaatat	7560
tactgggctg	ctattaacaa	gagatggtgg	taataacaac	aatgggtccg	agatcttcag	7620
acctggaggga	ggcgatatga	gggacaattg	gagaagtga	ttatataaat	ataaagtagt	7680
aaaaattgaa	ccattaggag	tagcaccac	caaggcaaag	agaagagtgg	tgacagagaga	7740
aaaaagagca	gtgggaatag	gagctttggt	ccttggttcc	ttgggagcag	caggaagcac	7800
tatgggctgc	acgtcaatga	cgctgacggt	acaggccaga	caattattgt	ctgatatagt	7860
gcagcagcag	aacaatttgc	tgagggtctat	tgaggcgcaa	cagcatctgt	tgcaactcac	7920
agtctggggc	atcaaacagc	tccaggcaag	aatcctggct	gtggaaagat	acctaaagga	7980
tcaacagctc	ctgggggattt	gggggtgtct	tggaaaactc	atttgcacca	ctgctgtgcc	8040
ttggaatgct	agttggagta	ataaatctct	ggaacagatt	tggaaataca	tgacctggat	8100
ggagtggggac	agagaaatta	acaattacac	aagcttaata	cactccttaa	ttgaagaatc	8160
gcaaaaccag	caagaaaaga	atgaacaaga	attattggaa	ttagataaat	gggcaagttt	8220
gtggaattgg	tttaacataa	caaattggct	gtggtatata	aaattattca	taatgatagt	8280
aggaggcttg	gtaggtttaa	gaatagtttt	tgctgtactt	tctatagtga	atagagttag	8340
gcagggatat	tcaccattat	cgtttcagac	ccacctccca	atcccagggg	gacccgacag	8400
gcccgaagga	atagaagaag	aaggtggaga	gagagacaga	gacagatcca	ttcgattagt	8460
gaacggatcc	ttagcactta	tctgggacga	tctgcgagac	ctgtgacctc	tcagctacca	8520
ccgcttgaga	gacttactct	tgattgttaac	gaggatttgt	gaacttctgg	gacgcagggg	8580
gtgggaagcc	ctcaaatatt	ggtggaaatct	cctacagtat	tggagtcagg	aactaaagaa	8640
tgtgtgctgt	aacttgctca	atgccacagc	catagcagta	gctgagggga	cagatagggt	8700
tatagaagta	ttacaagcag	cttatagagc	tattcgccac	atacctagaa	gaataagaca	8760

100T000"05T000

Ile	Ser	Thr	Ser	Ile	Arg	Asp	Lys	Val	Gln	Lys	Glu	Tyr	Ala	Phe	Phe	
				165					170					175		
Tyr	Lys	Leu	Asp	Ile	Val	Pro	Ile	Asp	Asn	Thr	Ser	Tyr	Arg	Leu	Ile	
			180					185					190			
Ser	Cys	Asn	Thr	Ser	Val	Ile	Thr	Gln	Ala	Cys	Pro	Lys	Val	Ser	Phe	
		195					200					205				
Glu	Pro	Ile	Pro	Ile	His	Tyr	Cys	Ala	Pro	Ala	Gly	Phe	Ala	Ile	Leu	
	210					215					220					
Lys	Cys	Asn	Asn	Lys	Thr	Phe	Asn	Gly	Thr	Gly	Pro	Cys	Thr	Asn	Val	
225					230					235					240	
Ser	Thr	Val	Gln	Cys	Thr	His	Gly	Ile	Arg	Pro	Val	Val	Ser	Thr	Gln	
				245					250						255	
Leu	Leu	Leu	Asn	Gly	Ser	Leu	Ala	Glu	Glu	Asp	Val	Val	Ile	Arg	Ser	
			260					265						270		
Ala	Asn	Phe	Thr	Asp	Asn	Ala	Lys	Thr	Ile	Ile	Val	Gln	Leu	Asn	Thr	
		275					280					285				
Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr	Arg	Lys	Ser	
		290					295				300					
Ile	Arg	Ile	Gln	Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Thr	Ile	Gly	Lys	
305					310					315					320	
Ile	Gly	Asn	Met	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Ala	Lys	Trp	
				325					330					335		
Asn	Ala	Thr	Leu	Lys	Gln	Ile	Ala	Ser	Lys	Leu	Arg	Glu	Gln	Phe	Gly	
			340					345					350			
Asn	Asn	Lys	Thr	Ile	Ile	Phe	Lys	Gln	Ser	Ser	Gly	Gly	Asp	Pro	Glu	
		355					360					365				
Ile	Val	Thr	His	Ser	Phe	Asn	Cys	Gly	Gly	Glu	Phe	Phe	Tyr	Cys	Asn	
	370					375					380					
Ser	Thr	Gln	Leu	Phe	Asn	Ser	Thr	Trp	Phe	Asn	Ser	Thr	Trp	Ser	Thr	
385					390					395					400	
Glu	Gly	Ser	Asn	Asn	Thr	Glu	Gly	Ser	Asp	Thr	Ile	Thr	Leu	Pro	Cys	
			405						410					415		
Arg	Ile	Lys	Gln	Phe	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met	
			420					425					430			
Tyr	Ala	Pro	Pro	Ile	Ser	Gly	Gln	Ile	Arg	Cys	Ser	Ser	Asn	Ile	Thr	
		435					440					445				
Gly	Leu	Leu	Leu	Thr	Arg	Asp	Gly	Gly	Asn	Asn	Asn	Asn	Gly	Ser	Glu	

460

Glu Arg Asp Arg Asp Arg Ser Ile Arg Leu Val Asn Gly Ser Leu Ala
740 745 750

[illegible]

Leu Ile Trp Asp Asp Leu Arg Ser Leu Cys Leu Phe Ser Tyr His Arg
755 760 765

Leu Arg Asp Leu Leu Leu Ile Val Thr Arg Ile Val Glu Leu Leu Gly
770 775 780

Arg Arg Gly Trp Glu Ala Leu Lys Tyr Trp Trp Asn Leu Leu Gln Tyr
785 790 795 800

Trp Ser Gln Glu Leu Lys Asn Ser Ala Val Asn Leu Leu Asn Ala Thr
805 810 815

Ala Ile Ala Val Ala Glu Gly Thr Asp Arg Val Ile Glu Val Leu Gln
820 825 830

Ala Ala Tyr Arg Ala Ile Arg His Ile Pro Arg Arg Ile Arg Gln Gly
835 840 845

Leu Glu Arg Ile Leu Leu
850

<210> 3

<211> 107

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of the artificial sequence:
oligonucleotide for cloning

<400> 3

aagatgtagt aattagatct gccaatattca cagacaatgc taaaaccata atagtagcagc 60
tgaacacatc gttagaaatt aattgtacaa gacccaacaa caataca 107

<210> 4

<211> 120

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of the artificial sequence:
oligonucleotide for cloning

<220>

<221> misc_feature

<222> (97)..(99)

<223> Sequence at this position: (GA)(AT)(GATC), ie.
base at position 97 can be G or A, base at
position 98 can be A or T, and base at
position 99 can be G, A, T or C.

<400> 4

ttttgtctta gaaatgttac aatgtgcttg tcttatgtct cctgttgcag cttctgttgc 60
atgaaatgct ctccctgggc cgatatggat actatgrwnt tttcttgtat tgttggttggg 120

FOOTNOTES: 667660

<210> 5
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of the artificial sequence:
sequencing primer

<400> 5
ccatgtacaa atgtcag

17

<210> 6
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of the artificial sequence:
sequencing primer

<400> 6
aaaactgtgc gttacaa

17

<210> 7
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of the artificial sequence:
sequencing primer

<400> 7
gtaaaacgac ggccagt

17

<210> 8
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of the artificial sequence:
sequencing primer

<400> 8
caggaaacag ctatgac

17

<210> 9
<211> 2148

<212> DNA
<213> Artificial Sequence

<220>

<223> Description of the artificial sequence: synthetic DNA

<220>

<221> misc_feature

<222> (3)..(9)

<223> BstEII cleavage site

<220>

<221> misc_feature

<222> (2143)..(2148)

<223> BamHI cleavage site

<400> 9

tgggtcaccg	tctattatgg	ggtgcctgtg	tggaaggaag	caaccaccac	tctattttgt	60
gcatcagatg	ctaaagcata	tgatacacag	gtacataatg	tttgggccac	acatgcctgt	120
gtaccacacg	acccaacccc	acaagaagta	gtattggtaa	atgtgacaga	aaattttaac	180
atgtggaaaa	atgacatggt	agaacagatg	catgaggata	taatcagttt	atgggatcaa	240
agccttaagc	catgtgtaaa	attaaccccc	ctctgtgtta	gtttaaagtg	cactgatttg	300
aagaatgata	ctaataccaa	tagtagtagc	gggagaatga	taatggagaa	aggagagata	360
aaaaactgca	gcttcaatat	cagcacaaagc	ataagagata	aggtgcagaa	agaatatgca	420
ttcttttata	aacttgatat	agtaccaata	gataatacca	gctatagggt	gataagttgt	480
aacacctcag	tgatcacaca	ggcctgtcca	aaggatcctt	ttgagccaat	tcccatacat	540
tattgtgccc	cggctggttt	tgcgattcta	aaatgtaata	ataagacgtt	caatggaaca	600
ggaccatgta	caaatgtcag	cacagtacaa	tgtacacatg	gaattcgacc	agtagtatca	660
actcaactgc	tgttaaatgg	cagtctagca	gaagaagatg	tagtaattag	atctgccaat	720
ttcacagaca	atgctaaaac	cataatagta	cagctgaaca	catctgtaga	aattaattgt	780
acaagacca	acaacaatac	aagaaaaagt	atccgtatcc	agaggggacc	aggagagaca	840
tttgttacaa	taggaaaaat	aggaaatatg	agacaagcac	attgtaacat	ttctagagca	900
aaatggaatg	ccacttttaa	acagatagct	agcaaattaa	gagaacaatt	tggaaataat	960
aaaacaataa	tctttaagca	gtcatccgga	ggggacccag	aaattgtaac	gcacagtttt	1020
aattgtggag	gggaattttt	ctactgtaat	tcaacacaac	tgtttaatat	tacttggttt	1080
aatagtactt	ggagtactga	aggggtcaat	aacactgaag	gaagtgcac	aatcacactc	1140
ccatgcagaa	taaaacaatt	tataaacatg	tggcaggaag	taggaaaagc	aatgtatgcc	1200
cctcccatca	gtggccaaat	tagatgttca	tcaaataatta	ctgggctgct	attaactcga	1260
gatggtggta	ataacaacaa	tgggtccgag	attttcagac	ctggaggagg	cgatatgagg	1320
gataattgga	gaagtgaatt	atataaatat	aaagtagtaa	aaattgaacc	attaggagta	1380
gcaccacca	aggcaaagag	acgcgtggtg	cagagagaaa	agcgcgcagt	gggaatagga	1440
gctctgttcc	ttgggttctt	gggagcagca	ggaagcacta	tgggcgcagc	gtcaatgacg	1500
ctgacggtac	aggccagaca	attattgtct	gatatagtgc	agcagcagaa	caatttgctg	1560
agggcaattg	aggcgcaaca	gcatctgttg	caactcacag	tctggggcat	caaacagctc	1620
caggcaagaa	tcctggctgt	ggaaagatac	ctaaaggatc	aacagctcct	ggggatttgg	1680
ggttgctctg	gaaaactcat	ttgcaccact	gctgtgcctt	ggaatgctag	ttggagtaat	1740
aaatctctgg	aacagatttg	gaataacatg	acctggatgg	agtgggacag	agaaattaac	1800
aattacacaa	gcttaataca	ctccttaatt	gaagaatcgc	aaaaccagca	agaaaagaat	1860
gaacaagaat	tattggaatt	agataaatgg	gcaagtttgt	ggaattgggt	taacataaca	1920
aattggctgt	ggtatataaa	attattcata	atgatagtag	gaggcttggt	aggtttaaga	1980
atagtttttg	ctgtactttc	tatagtgaat	agagttaggc	agggatattc	accattatcg	2040
tttcagaccc	acctcccaat	cccaggggga	cccagacaggc	ccgaaggaat	agaagaagaa	2100
ggtggagaga	gagacagaga	cagatccatt	cgattagtga	acggatcc		2148

<210> 10

<211> 6229

ccacacatgc	ctgtgtacc	acagacccca	accacaaga	agtagtattg	gtaaatgtga	1560
cagaaaat	taacatgtg	aaaaatgaca	tggtagaaca	gatgcattgag	gatataatca	1620
gtttatggga	tcaaagccta	aagccatgtg	taaaattaac	cccactctgt	gttagtttaa	1680
agtgcactga	tttgaagaat	gatactaata	ccaatagtag	tagcgggaga	atgataatgg	1740
agaaaggaga	gataaaaaac	tgctctttca	atatcagcac	aagcataaga	gataaggtgc	1800
agaaagaata	tgcattcttt	tataaacttg	atatagtacc	aatagataat	accagctata	1860
ggttgataag	ttgtaacacc	tcagtcatta	cacaggcctg	tccaaaggta	tcctttgagc	1920
caattcccat	acattattgt	gcccggctg	gttttgcgat	tctaaaatgt	aataataaga	1980
cgttcaatgg	aacaggacca	tgtacaaatg	tcagcacagt	acaatgtaca	catggaatca	2040
ggccagtagt	atcaactcaa	ctgctgttaa	atggcagctc	agcagaagaa	gatgtagtaa	2100
ttagatctgc	caatttcaca	gacaatgcta	aaaccataat	agtacagctg	aacacatctg	2160
tagaaattaa	ttgtacaaga	cccaacaaca	atacaagaaa	aagtatccgt	atccagaggg	2220
gaccagggag	agcatttggt	acaataggaa	aaataggaaa	tatgagacaa	gcacattgta	2280
acattagtag	agcaaaaatg	aatgccactt	taaaacagat	agctagcaaa	ttaagagAAC	2340
aatttggaAA	taataaaaaca	ataatcttta	agcaatcctc	aggagggggac	ccagaaattg	2400
taacgcacag	ttttaattgt	ggaggggaat	ttttctactg	taattcaaca	caactgttta	2460
atagtacttg	gtttaatagt	acttgaggta	ctgaagggtc	aaataacact	gaagggaagt	2520
acacaatcac	actcccatgc	agaataaaaac	aattttataaa	catgtggcag	gaagtaggaa	2580
aagcaatgta	tgcctctccc	atcagtggac	aaattagatg	ttcatcaaat	attactgggc	2640
tgctattaac	aagagatgg	ggtaataaca	acaatgggtc	cgagatcttc	agacctggag	2700
gaggcgatat	gagggacaat	tggaagagtg	aattatataa	atataaagta	gtaaaaattg	2760
aaccattagg	agtagcaccc	accaaggcaa	agagaagagt	ggtgcagaga	gtaaaaagag	2820
cagtgggaat	aggagctttg	ttccttggg	tcttgggagc	agcaggaagc	actatgggct	2880
gcacgtcaat	gacgctgacg	gtacaggcca	gacaattatt	gtctgatata	gtgcagcagc	2940
agaacaat	gctgagggct	attgaggcgc	aacagcatct	gttgcaactc	acagtctggg	3000
gcatcaaaca	gctccaggca	agaatcctgg	ctgtggaaag	atacctaag	gatcaacagc	3060
tcctggggat	ttgggggttg	tctggaaaac	tcatttgcac	cactgctgtg	ccttggaatg	3120
ctagttagg	taataaatct	ctggaacaga	tttggaataa	catgacctgg	atggagtggg	3180
acagagaaat	taacaattac	acaagcttaa	tacactcctt	aattgaagaa	tcgcaaaacc	3240
agcaagaaaa	gaatgaacaa	gaattattgg	aattagataa	atgggcaagt	ttgtggaatt	3300
ggtttaacat	aacaaattgg	ctgtggtata	taaaattatt	cataatgata	gtaggaggct	3360
tggtaggttt	aagaatagtt	tttgctgtac	tttctatagt	gaatagagtt	aggcagggat	3420
attcaccatt	atcgtttcag	accacctcc	caatcccag	gggacccgac	aggcccgaag	3480
gaatagaaga	agaagggtga	gagagagaca	gagacagatc	cattcgatta	gtgaacggat	3540
ccttagcact	tatctgggac	gatctgcgga	gcctgtgcct	cttcagctac	caccgcttga	3600
gagacttact	cttgattgta	acgaggattg	tggaacttct	gggacgcagg	gggtgggaag	3660
ccctcaaata	ttggtggaat	ctcctacagt	attggagtca	ggaactaaag	aatagtgtct	3720
ttaaacttgct	caatgccaca	gccatagcag	tagctgaggg	gacagatagg	gttataaga	3780
tattacaagc	agcttataga	gctattcgcc	acatacctag	aagaataaga	cagggtcttg	3840
aaaggatttt	gctataagat	gggtggcaag	tggtcaaaaa	gtagtgtgat	tggtatggct	3900
gctgtaaggg	aaagaatgag	acgagctgag	ccagcagcag	atgggggtgg	agcagtatct	3960
cgagatctag	actagaacta	gcttcgatcc	agacatgata	agatacattg	atgagttttg	4020
acaaaccaca	actagaatgc	agtgaaaaaa	atgctttatt	tgtgaaattt	gtgatgctat	4080
tgctttat	gtaaccatta	taagctgcaa	taaacaagtt	aacaacaaca	attgcattca	4140
ttttatgttt	caggttcagg	gggaggtgtg	ggaggttttt	taaagcaagt	aaaacctcta	4200
caaatgtggt	atggctgatt	atgatcctgc	ctcgcgcggt	tcggtgatga	cggtgaaaac	4260
ctctgacaca	tgcagctccc	ggagacggtc	acagcttgtc	tgtaagcgga	tgccggggagc	4320
agacaagccc	gtcagggcgc	gtcagcgggt	gttgccgggt	gtcggggcgc	agccatgacc	4380
cagtcaagta	gcgatagcgg	agtgtatact	ggcttaacta	tgccgcatca	gagcagattg	4440
tactgagagt	gcaccatatg	tcgggcgcgc	ttgctggcgt	ttttccatag	gctccgcccc	4500
cctgacgagc	atcacaaaaa	tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	4560
taaagatacc	aggcgtttcc	ccctggaagc	tcctctgtgc	gctctcctgt	tcggacctgt	4620
ccgcttaccg	gataacctgc	cgcctttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	4680
tcacgctgta	ggtatctcag	ttcggtgtag	gtcgttcgct	ccaagctggt	ctgtgtgcac	4740
gaaccccccg	ttcagcccca	cgcgtgcgcc	ttatccggta	acatcgtct	tgagtccaac	4800
ccggtaaagac	acgacttatc	gccactggca	gcagccactg	gtaacaggat	tagcagagcg	4860
aggatatgtag	gcggtgctac	agagttcttg	aagtgggtggc	ctaactacgg	ctacactaga	4920

```

aggacagtat ttggtatctg cgctctgctg aagccagtta ccttcggaaa aagagttggt 4980
agctcttgat ccggaaca aaccaccgct ggtagcgggtg gttttttgt ttgcaagcag 5040
cagattacgc gcagaaaaaa aggatctcaa gaagatcctt tgatcttttc tacgggggtct 5100
gacgctcagt ggaacgaaaa ctcacgttaa gggatttttg tcatgagatt atcaaaaagg 5160
atcttcacct agatcctttt aaattaaaaa tgaagtttta aatcaatcta aagtatatat 5220
gagtaaactt ggtctgacag ttaccaatgc ttaatcagtg aggcacctat ctcagcgatc 5280
tgtctatttc gttcatccat agttgctga ctccccgctg tgtagataac tacgatacgg 5340
gagggcttac catctggccc cagtgtgtga atgataccgc gagaccacg ctcaccgggt 5400
ccagatttat cagcaataaa ccagccagcc ggaagggccg agcgcagaag tggtcctgca 5460
actttatccg cctccatcca gtctattaat tgttgccggg aagctagagt aagtagttcg 5520
ccagttaata gtttgcgcaa cgttgttgcc attgtctacag gcatcgtggt gtcacgctcg 5580
tcgtttggta tggcttcatt cagctccggt tcccaacgat caaggcgagt tacatgatcc 5640
cccatgttgt gcaaaaaagc ggttagctcc ttcggtcctc cgatcgttgt cagaagtaag 5700
ttggccgcag tggtatcact catggttatg gcagcactgc ataattctct tactgtcatg 5760
ccatccgtaa gatgcttttc tgtgactggt gagtactcaa ccaagtcatt ctgagaatag 5820
tgtatgcggc gaccgagttg ctcttgccc gcgtaatac gggataatac cgcgccacat 5880
agcagaactt taaaagtgtc catcattgga aaacgttctt cggggcgaaa actctcaagg 5940
atcttaccgc tgttgagatc cagttcgatg taaccactc gtgcaccaa ctgatcttca 6000
gcatctttta ctttcaccag cgtttctggg tgagcaaaaa caggaaggca aaatgccgca 6060
aaaaagggaa taagggcgac acggaaatgt tgaatactca tactcttctt ttttcaatat 6120
tattgaagca tttatcaggg ttattgtctc atgagcggat acatatttga atgtatttag 6180
aaaaataaac aaataggggt tccgcgcaca tttccccgaa aagtgccac 6229

```

```

<210> 11
<211> 860
<212> DNA
<213> Human immunodeficiency virus

<220>
<221> misc_feature
<222> (1)..(860)
<223> PI-932 original sequence V1-V2-V3-loop

```

```

<400> 11
tgtgtaccca cagaccccaa cccacaaaag gtagtattgg aaaatgtgac agaaaatttt 60
aacatgtgga aaaatgacat ggtagaacag atgcatgagg atataatcaa tttatgggat 120
caaagcctaa agccatgtgt aaaactaacc ccactctgtg ttacttttaa ttgcactgat 180
gctgatttaa attgcaataa tactgattta aattgcacta aagctaattt ggggaaaaat 240
actcataaca atactattag tgggaaaata atagagaag tagaaataaa aaactgctct 300
ttcaagggtca ccacaggcat aagggtataag atgcaaaaag aatatgcact tttgaataaa 360
cttgatatag taccaataga taatgataag aataatacta actttatatt gataagttgt 420
aacacctcga ccattacaca ggctgtgcca aaggatatct ttgagccaat tccatacat 480
ttttgtgccc cggtgtggtt tgcgattcta aagtgtaatg aaaagagtta cagtggaaaa 540
ggaccatgta aaaatgtcag cacagtacaa tgtacacatg gaattaggcc agtagtgtca 600
actcaactgc tgttgaatgg cagtctagca gaaaaagaag tagtaattag atctgagaat 660
ttcacagaca atgctaaaac cataatagta cagctgaagg aatctgtaaa cattacttgt 720
ataagacccc acaacactgt aacagacagg atacatatag ggccagggag atcatttcat 780
acaacaagaa aaataaaagg agatataaga caagcacatt gtagccttag gagaaaagat 840
tggataaaca ctttacaaga
860

```

```

<210> 12
<211> 870
<212> DNA
<213> Artificial Sequence

```

[illegible]

<400> 12

tgtgtacc	caagccca	cccacaaa	gtagtatt	aaaatgtg	agaaaaat	60
aatatgtg	aaaatgac	ggtagaac	atgcatga	atataatc	tttatggg	120
caaagcct	agccatgt	aaaactaa	ccactctg	ttacttta	ttgactga	180
gctgattta	attgcaata	tactgatt	aattgcac	aagctaatt	ggggaaaa	240
actcatact	gcagtatt	tgggaaa	atagagaa	tagaaata	aaactgct	300
ttcaagggt	ccacaggc	aagggata	atgcaaaa	aatatgca	tttgaata	360
cttgatat	taccaata	taatgata	aataatac	actttata	gataagtt	420
aacacctcg	tgatcac	ggcctgtc	aaggatcc	ttgagcca	tcccatac	480
ttttgtgcc	cggtgtgt	tgcgattc	aagtgtat	aaaagagt	cagtggaa	540
ggaccatgt	aaaatgtc	cacagtac	tgtacac	gaattcgg	agtagtgt	600
actcaactg	tgttgaat	cagtctag	gaaaaaga	tagtaatt	atctgaga	660
ttcacagac	atgtcaaaa	cataatag	cagctga	aatctgt	cattactt	720
ataagacccc	acaactact	aacagacg	atacatat	ggccaggg	atcatttc	780
acaacaaga	aaataaaa	agatataa	caagcac	gtagcctt	tagaaaag	840
tggaataac	ctttaca	gatagcta				870

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

SCHREIBER

Atty. Ref.: 35-213

Serial No. Unknown

Group:

U.S. National Phase of PCT/EP99/09759

Filed: August 10, 2001

Examiner:

For: VIRAL VACCINE

* * * * *

August 10, 2001

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

LETTER


The attached paper and computer readable copies of the Sequence Listing are the same.

No new matter has been added.

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: _____



B. J. Sadoff
Reg. No. 36,663

1100 North Glebe Road, 8th Floor
Arlington, VA 22201-4714
Telephone: (703) 816-4000
Facsimile: (703) 816-4100

SEQUENCE LISTING

<110> Strathmann AG & Co.

<120> Virus-Vaccine

<130> P057760

<140>

<141>

<150> 199 07 485.2

<151> 1999-02-12

<160> 12

<170> PatentIn Ver. 2.1

<210> 1

<211> 9709

<212> DNA

<213> Human immunodeficiency virus

<400> 1

tggaagggct	aatttgggtc	caaaaaagac	aagagatcct	tgatctgtgg	atctaccaca	60
caCaaggcta	cttccctgat	tggcagaact	acacaccagg	gccagggatc	agatatccac	120
tgacctttgg	atgggtgcttc	aagtttagtac	cagttgaacc	agagcaagta	gaagaggcca	180
aataaggaga	gaagaacagc	ttgttacacc	ctatgagcca	gcatgggatg	gaggaccgg	240
agggagaagt	attagtgtgg	aagtttgaca	gcctcctagc	atttcgtcac	atggcccag	300
agctgcaccc	ggagtactac	aaagactgct	gacatcgagc	tttctacaag	ggactttccg	360
ctggggactt	tccagggagg	tgtggcctgg	gcgggactgg	ggagtggcga	gccctcagat	420
gctacatata	agcagctgct	ttttgcctgt	actgggtctc	tctgggttaga	ccagatctga	480
gcctgggagc	tctctgggcta	actaggggaa	ccactgctta	agcctcaata	aagcttgcct	540
tgagtgtctca	aagtagtgtg	tgcccgctctg	ttgtgtgact	ctggtaacta	gagatccctc	600
agaccctttt	agtcagtgtg	gaaaatctct	agcagtggcg	cccgaacagg	gacttgaaag	660
cgaaagtata	gccagaggag	atctctcgac	gcaggactcg	gcttgctgaa	gcgcgcacgg	720
caagaggcga	ggggcggcga	ctgggtgagta	cgccaaaaat	tttgactagc	ggaggctaga	780
aggagagaga	tgggtgcgag	agcgtcggta	ttaagcgggg	gagaattaga	taaatgggaa	840
aaaaattcgg	taaggccagg	gggaaagaaa	caatataaac	taaaacatat	agtatgggca	900
agcaggggagc	tagaacgatt	cgcagttaat	cctggccttt	tagagacatc	agaaggctgt	960
agacaaatac	tgggacagct	acaaccatcc	cttcagacag	gatcagaaga	acttagatca	1020
ttatataata	caatagcagt	cctctattgt	gtgcatcaaa	ggatagatgt	aaaagacacc	1080
aaggaaggcct	tagataagat	agaggaagag	caaaacaaaa	gtaagaaaaa	ggcacagcaa	1140
gcagcagctg	acacaggaaa	caacagccag	gtcagccaaa	attaccctat	agtgcagaac	1200
ctccaggggc	aaatggtaca	tcaggccata	tcacctagaa	ctttaaatgc	atgggtaaaa	1260
gtagtagaag	agaaggcttt	cagcccagaa	gtaataccca	tgttttcagc	attatcagaa	1320
ggagccaccc	cacaagattt	aaataccatg	ctaaacacag	tggggggaca	tcaagcagcc	1380
atgcaaatgt	taaaagagac	catcaatgag	gaagctgcag	aatgggatag	attgcattca	1440
gtgcatgcag	ggcctattgc	accaggccag	atgagagaac	caagggggaag	tgacatagca	1500
ggaactacta	gtacccttca	ggaacaaaata	ggatggatga	cacataatcc	acctatccca	1560
gtaggagaaa	tctataaaaag	atggataatc	ctgggattaa	ataaaatagt	aagaatgtat	1620
agccctacca	gcattcttga	cataagacaa	ggaccaaagg	aacccttttag	agactatgta	1680
gaccgattct	ataaaaactct	aagagccgag	caagcttcac	aagaggtaaa	aaatttggatg	1740
acagaaacct	tgttgggtcca	aaatgcgaac	ccagatttga	agactatttt	aaaagcattg	1800
ggaccaggag	cgacactaga	agaaatgatg	acagcatgtc	agggagtggg	gggacccggc	1860
cataaaggcaa	gagtttttggc	tgaagcaatg	agccaagtaa	caaatccagc	taccataatg	1920
atacagaaga	gcaatttttag	gaaccaaaga	aagactgtta	agtgttttcaa	ttgtggcaca	1980
gaagggcaca	tagccaaaaa	ttgcaggggcc	cctaggaaaa	agggctgttg	gaaatgtgga	2040
aaggaaggac	accaaatagaa	agattgtact	gagagacagg	ctaatttttt	aggggaagatc	2100
tggccttccc	acaaggggaag	gccagggaat	tttcttcaga	gcagaccaga	gccaacagcc	2160
ccaccagaag	agagcttcag	gtttggggaa	gagacaacaa	ctccctctca	gaagcaggag	2220
ccgatagaca	aggaactgta	tccttttagct	tccttcagat	cactcttttg	cagcgacctc	2280
tcgtcacaa	aaagataggg	gggcaattaa	aggaagctct	attagatata	ggagcagatg	2340
atacagtatt	agaagaaatg	aatttgcag	gaagatggaa	acaaaaaatg	atagggggaa	2400
ttggagggtt	tatcaaagta	ggacagtatg	atcagatact	catagaaatc	tgcggacata	2460
aagctatagg	tacagtatta	gtaggacctc	ccactgtcaa	cataattgga	agaaatctgt	2520
tgactcagat	tggctgcact	ttaaattttc	ccattagtcc	tattgagact	gtaccagtaa	2580
aattaaagcc	aggaatggat	ggcccaaaa	ttaaacaatg	gccattgaca	gaagaaaaaa	2640

09/913159-518-1

[illegible]

gataatggag	aaaggagaga	taaaaaactg	ctcttttcaat	atcagcacaa	gcataagaga	6720
taaggtgcag	aaagaatatg	cattctttta	taaacttgat	atagtagcaa	tagataatac	6780
cagctatagg	ttgataagtt	gtaaacacct	agtcattaca	caggcctgtc	caaagggtatc	6840
ctttgagcca	attcccatac	attattgtgc	ccggctgtgt	tttgcgattc	taaaatgtaa	6900
taataagacg	ttcaatggaa	caggaccatg	tacaaatgtc	agcagcgtac	aatgtacaga	6960
tggaatcagg	ccagtagtat	caactcaact	gctgttaa	ggcagcttag	cagaagacaa	7020
tgtagttaatt	agatctgcca	attttcacaga	caatgctaaa	accataatag	tacagctgaa	7080
cacatctgta	gaaatatta	gtacaagacc	caacaacaa	acaagaaaaa	gtatccgtat	7140
ccagagggga	ccagggaagt	catttgttac	aataggaaaa	ataggaaata	tgagacaagc	7200
acattgttaac	attagtagag	caaaatggaa	tgccacttta	aaacagatag	ctagcaaat	7260
aagagaacaa	tttggaata	ataaaacaat	aatctttaag	caatcctcag	gaggggaccc	7320
agaaatgtta	acgcacagtt	ttaattgtgg	aggggaattt	ttctactgta	attcaacaca	7380
actgtttta	agtacttggt	ttaatagtc	ttggagtact	gaagggtcaa	ataacactga	7440
aggaagtgc	acaatcacac	tcccatgcag	aataaaacaa	ttataaaca	tgtggcagga	7500
agtaggaaaa	gcaatgtatg	ccctcccat	cagtggacaa	attagatggt	catcaaat	7560
tactgggctg	ctattaacaa	gagatggtgg	taataacaac	aatgggtccg	agatcttcag	7620
acctggagga	ggcgatatga	gggacaattg	gagaagtga	ttatataaat	ataaagtagt	7680
aaaaattgaa	ccctatggag	tagcaccac	caaggcgaag	agaagagtgg	tcgagagaga	7740
aaaaagagca	gtgggaatag	gagctttgtt	ccttgggttc	ttgggagcag	caggaagcac	7800
tatgggctgc	acgtcaatga	cgctgacggt	acaggccaga	caattattgt	ctgatatagt	7860
gcagcagcag	aacaatttgc	tgagggtcat	tgaggcgcaa	cagcatctgt	tgcaactcac	7920
agtctggggc	atcaaacagc	tccaggcaag	aatcctggct	gtggaaagat	acctaaagga	7980
tcaacagctc	ctggggattt	gggttgcctc	tggaaaactc	atttgcacca	ctgctgtgcc	8040
ttggaatgct	agttggagta	ataaatctct	ggaacagatt	tggaaataaca	tgacctggat	8100
ggagtgggac	agagaaatta	acaattacac	aagcttaata	cactccttaa	ttgaagaatc	8160
gcaaaactcg	caagaaaaa	atgaacaaga	attattggaa	ttagataaat	gggcaagttt	8220
gtggaattgg	tttaacataa	caaattggct	gtggtatata	aaattattca	taatgattag	8280
aggaggcttg	ttagggttaa	gaatagtttt	tgctgtactt	cttatagtga	atagagttag	8340
gcagggatat	tcaccattat	cgtttcagac	ccactccca	atcccgaggg	gaccgcagac	8400
gcccgaagga	atagaagaag	aaggtggaga	gagagacaga	gacagatcca	ttcgattagt	8460
gaacgtagtc	ttagcactta	tctgggacga	tctgaggagc	ctgtgcctct	tcagctacca	8520
ccgcttgaga	gacttactct	tgtattgaac	gaggatttgt	gaactcttgg	gacgcagggg	8580
gtgggaagcc	ctcaaatatt	ggtggaaatc	cctacagtat	tggagtccag	aactaaagaa	8640
tagtgctggt	aacttgctca	atgccacagc	catagcagta	gctgagggga	cagatagggt	8700
tatagaagta	ttacaagcag	cttatagagc	tattcgccac	atacctagaa	gaataagaca	8760
gggcttgga	aggattttgc	tataagattg	gtggcaagtg	gtcaaaaagt	agtgtgattg	8820
tatggccttc	tgtaaaggaa	agaatgagac	gagctagagc	agcagcagat	gggggtggag	8880
cagtatctcg	agacctagaa	aaacatggag	caatcacaa	tagcaataca	gcagctaaaca	8940
atgctgcttg	tgcttggtca	gaagcacaa	aggaggaaga	ggtgggtttt	ccagtcacac	9000
ctcaggtacc	tttaagacca	atgacttaca	aggcagctgt	agatcttagc	cactttttaa	9060
aagaaaaggg	gggactggaa	gggctaattc	actcccaag	aagacaagat	atccttgatc	9120
tgtggatcta	ccacacacaa	ggctacttcc	ctgattggca	gaactacaca	ccaggggccag	9180
gggtcagata	tccactgacc	tttggatggt	gctacaagct	agtaccagtt	gagccagata	9240
aggtagaaga	ggccaataaa	ggagagaaca	ccagcttgtt	acaccctgtg	agcctgcattg	9300
gaatggatga	ccctgagaga	gaagtgttag	agtggagggt	tgacagccgc	ctagcatttc	9360
atcacgtggc	ccgagagctg	catccggagt	acttcaagaa	ctgctgacat	cgagcttggt	9420
acaagggact	ttccgctggg	gactttccag	ggaggcgtgg	cctggggcggg	actggggagt	9480
ggcgagccct	cagatgctgc	atataagcag	ctgctttttg	cctgtactgg	gtctctctgg	9540
ttagaccaga	cttgagcctg	ggagctctct	ggctaactag	ggaacccact	gcttaagcct	9600
caataaagct	tgcccttgagt	gcttcaagta	gtgtgtccc	gtctgttgtg	tgactctggt	9660
aactagagat	ccctcagacc	cttttagtca	gtgtggaaaa	tctctagca		9709

<220>
<223> Envelope Polyprotein

```

<400> 2
Met Arg Val Lys Glu Lys Tyr Gln His Leu Trp Arg Trp Gly Trp Lys
  1      5      10      15
Trp Gly Thr Met Leu Leu Gly Ile Leu Met Ile Cys Ser Ala Thr Glu
      20      25      30
Lys Leu Trp Val Thr Val Tyr Tyr Gly Val Pro Val Trp Lys Glu Ala
      35      40      45
Thr Thr Thr Leu Phe Cys Ala Ser Asp Ala Lys Ala Tyr Asp Thr Glu
      50      55      60
Val His Asn Val Trp Ala Thr His Ala Cys Val Pro Thr Asp Pro Asn
      65      70      75      80
Pro Gln Glu Val Val Leu Val Asn Val Thr Glu Asn Phe Asn Met Trp
      85      90      95
Lys Asn Asp Met Val Glu Gln Met His Glu Asp Ile Ile Ser Leu Trp
      100      105      110
Asp Gln Ser Leu Lys Pro Cys Val Lys Leu Thr Pro Leu Cys Val Ser
      115      120      125
Leu Lys Cys Thr Asp Leu Lys Asn Asp Thr Asn Thr Asn Ser Ser Ser
      130      135      140
Gly Arg Met Ile Met Glu Lys Gly Glu Ile Lys Asn Cys Ser Phe Asn
      145      150      155      160
Ile Ser Thr Ser Ile Arg Asp Lys Val Gln Lys Glu Tyr Ala Phe Phe
      165      170      175
Tyr Lys Leu Asp Ile Val Pro Ile Asp Asn Thr Ser Tyr Arg Leu Ile
      180      185      190
Ser Cys Asn Thr Ser Val Ile Thr Gln Ala Cys Pro Lys Val Ser Phe
      195      200      205
Glu Pro Ile Pro Ile His Tyr Cys Ala Pro Ala Gly Phe Ala Ile Leu
      210      215      220
Lys Cys Asn Asn Lys Thr Phe Asn Gly Thr Gly Pro Cys Thr Asn Val
      225      230      235      240
Ser Thr Val Gln Cys Thr His Gly Ile Arg Pro Val Val Ser Thr Gln
      245      250      255
Leu Leu Leu Asn Gly Ser Leu Ala Glu Glu Asp Val Val Ile Arg Ser
      260      265      270
Ala Asn Phe Thr Asp Asn Ala Lys Thr Ile Ile Val Gln Leu Asn Thr
      275      280      285

```

Ser	Val	Glu	Ile	Asn	Cys	Thr	Arg	Pro	Asn	Asn	Asn	Thr	Arg	Lys	Ser
290						295					300				
Ile	Arg	Ile	Gln	Arg	Gly	Pro	Gly	Arg	Ala	Phe	Val	Thr	Ile	Gly	Lys
305					310					315					320
Ile	Gly	Asn	Met	Arg	Gln	Ala	His	Cys	Asn	Ile	Ser	Arg	Ala	Lys	Trp
				325					330					335	
Asn	Ala	Thr	Leu	Lys	Gln	Ile	Ala	Ser	Lys	Leu	Arg	Glu	Gln	Phe	Gly
			340					345					350		
Asn	Asn	Lys	Thr	Ile	Ile	Phe	Lys	Gln	Ser	Ser	Gly	Gly	Asp	Pro	Glu
		355					360					365			
Ile	Val	Thr	His	Ser	Phe	Asn	Cys	Gly	Gly	Glu	Phe	Phe	Tyr	Cys	Asn
	370					375					380				
Ser	Thr	Gln	Leu	Phe	Asn	Ser	Thr	Trp	Phe	Asn	Ser	Thr	Trp	Ser	Thr
385					390					395					400
Glu	Gly	Ser	Asn	Asn	Thr	Glu	Gly	Ser	Asp	Thr	Ile	Thr	Leu	Pro	Cys
				405					410					415	
Arg	Ile	Lys	Gln	Phe	Ile	Asn	Met	Trp	Gln	Glu	Val	Gly	Lys	Ala	Met
			420					425					430		
Tyr	Ala	Pro	Pro	Ile	Ser	Gly	Gln	Ile	Arg	Cys	Ser	Ser	Asn	Ile	Thr
		435					440					445			
Gly	Leu	Leu	Leu	Thr	Arg	Asp	Gly	Gly	Asn	Asn	Asn	Asn	Gly	Ser	Glu
	450					455					460				
Ile	Phe	Arg	Pro	Gly	Gly	Gly	Asp	Met	Arg	Asp	Asn	Trp	Arg	Ser	Glu
465					470					475					480
Leu	Tyr	Lys	Tyr	Lys	Val	Val	Lys	Ile	Glu	Pro	Leu	Gly	Val	Ala	Pro
				485					490					495	
Thr	Lys	Ala	Lys	Arg	Arg	Val	Val	Gln	Arg	Glu	Lys	Arg	Ala	Val	Gly
			500					505					510		
Ile	Gly	Ala	Leu	Phe	Leu	Gly	Phe	Leu	Gly	Ala	Ala	Gly	Ser	Thr	Met
		515					520					525			
Gly	Cys	Thr	Ser	Met	Thr	Leu	Thr	Val	Gln	Ala	Arg	Gln	Leu	Leu	Ser
	530					535					540				
Asp	Ile	Val	Gln	Gln	Gln	Asn	Asn	Leu	Leu	Arg	Ala	Ile	Glu	Ala	Gln
545					550					555					560
Gln	His	Leu	Leu	Gln	Leu	Thr	Val	Trp	Gly	Ile	Lys	Gln	Leu	Gln	Ala
				565					570					575	
Arg	Ile	Leu	Ala	Val	Glu	Arg	Tyr	Leu	Lys	Asp	Gln	Gln	Leu	Leu	Gly
			580					585					590		
Ile	Trp	Gly	Cys	Ser	Gly	Lys	Leu	Ile	Cys	Thr	Thr	Ala	Val	Pro	Trp
		595					600					605			
Asn	Ala	Ser	Trp	Ser	Asn	Lys	Ser	Leu	Glu	Gln	Ile	Trp	Asn	Asn	Met
	610					615					620				

Thr Trp Met Glu Trp Asp Arg Glu Ile Asn Asn Tyr Thr Ser Leu Ile
625 630 635 640

His Ser Leu Ile Glu Glu Ser Gln Asn Gln Gln Glu Lys Asn Glu Gln
645 650 655

Glu Leu Leu Glu Leu Asp Lys Trp Ala Ser Leu Trp Asn Trp Phe Asn
660 665 670

Ile Thr Asn Trp Leu Trp Tyr Ile Lys Leu Phe Ile Met Ile Val Gly
675 680 685

Gly Leu Val Gly Leu Arg Ile Val Phe Ala Val Leu Ser Ile Val Asn
690 695 700

Arg Val Arg Gln Gly Tyr Ser Pro Leu Ser Phe Gln Thr His Leu Pro
705 710 715 720

Ile Pro Arg Gly Pro Asp Arg Pro Glu Gly Ile Glu Glu Glu Gly Gly
725 730 735

Glu Arg Asp Arg Asp Arg Ser Ile Arg Leu Val Asn Gly Ser Leu Ala
740 745 750

Leu Ile Trp Asp Asp Leu Arg Ser Leu Cys Leu Phe Ser Tyr His Arg
755 760 765

Leu Arg Asp Leu Leu Leu Ile Val Thr Arg Ile Val Glu Leu Leu Gly
770 775 780

Arg Arg Gly Trp Glu Ala Leu Lys Tyr Trp Trp Asn Leu Leu Gln Tyr
785 790 795 800

Trp Ser Gln Glu Leu Lys Asn Ser Ala Val Asn Leu Leu Asn Ala Thr
805 810 815

Ala Ile Ala Val Ala Glu Gly Thr Asp Arg Val Ile Glu Val Leu Gln
820 825 830

Ala Ala Tyr Arg Ala Ile Arg His Ile Pro Arg Arg Ile Arg Gln Gly
835 840 845

Leu Glu Arg Ile Leu Leu
850

<210> 3
<211> 107
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of the artificial sequence:
oligonucleotide for cloning

<400> 3
aagatgtagt aattagatct gccaatctca cagacaatgc taaaaccata atagtacagc 60
tgaacacatc gttagaaatt aattgtacaa gacccaacaa caataca 107

```
<400> 7
gtaaaacgac ggccagt
```

17

<400> 9						
tgggtcaccg	tctattatgg	ggtgcctgtg	tggaaggaag	caaccaccac	tctattttgt	60
gcatcagatg	ctaaagcata	tgatacagag	gtacataatg	tttggggccac	acatgcctgt	120
gtacccacag	accccaatccc	acaagaagta	gtatttggtaa	atgtgcagaca	aaattttaac	180
atgtgtgaaaa	atgatcatggt	agaacagatg	catgaggata	taatcagcttt	atgggatcaa	240
agcctttaagc	catgtgtataa	attaacccca	ctctgtgttta	gtttaaagtg	cactgatttg	300
aagaatgata	ctaataccaa	tagtagtagc	gggagaatga	taatggagaa	aggagagata	360
aaaaactgca	gcttcaatat	cagcacaagc	ataagagata	agggtgcagaa	agaatatgca	420
ttcttttata	aacttगतat	agtaccaata	gataatacca	gctataggtt	gataagttgt	480
aacacctcag	tgatcacaca	ggcctgtcca	aaggtatcct	ttgagccaat	tcccatacat	540
tattgtgccc	cggtctggtt	tgcgattcta	aaatgttaata	ataagacgtt	caattggaaca	600
ggaccatgta	caaatgtcag	cacagtacaa	tgtacacatg	gaattcgacc	agtagtatca	660
actcaactgc	tgttaaatgg	cagtctagca	gaagaagatg	tagtaattag	atctgccaat	720
ttcacagaca	atgctaaaaa	cataatatga	cagctgaaca	catctgtaga	aattaattgt	780
acaagaccaca	acaactaatc	aagaaaaagt	atccgtatcc	agagggggacc	aggagagaca	840
tttgttacaa	taggaaaaat	aggaaatatg	agacaagcac	attgttaact	ttctagagca	900
aaatggaatg	ccacttttaa	acagatagct	agcaaattaa	gagaacaatt	tggaaataat	960
aaaacaataa	tctttaagca	gtcatccgga	ggggaccacc	aaatttgtaac	gcacagtttt	1020
aattgtggag	gggaattttt	ctactgtaat	tcaacacaa	tgttttaatat	tacttggttt	1080
aatagtactt	ggagtagctg	agggtcaaat	aacactgaag	gaagtgcac	aatcacactc	1140
ccatgcagaa	taaaacaatt	tataaacatg	tggcaggaa	taggaaaagc	aatgtatgcc	1200
cctcccatca	gtggccaaat	tagatgttca	tcaaatatta	ctgggctgct	attaactcga	1260
gatgggtggta	ataacaacaa	tgggtccgag	attttcagac	ctggaggagg	cgatatgagg	1320
gataatttga	gaagtgaatt	atataaatat	aaagtagtaa	aaattgaacc	attaggagta	1380
gcacccacca	aggcaaaggag	acgcgtggta	cagagagaaa	agcgcgcagt	gggaatatga	1440
gtctctgttc	ttgggttctt	gggagcagca	ggaagcacta	tgggcgcagc	gtcaatgcag	1500
ctgacgggtac	aggccagaca	attattgtct	gatatatgtc	agcagcagaa	caatttgcctg	1560
agggcaattg	aggcgcaaca	gcattctgtt	caactcacag	tctggggcat	caaacagctc	1620
caggcaagaa	tcttggtctg	ggaaagatac	ctaaaggtac	aacagctcct	ggggatttgg	1680
gggtgtctctg	gaaaactcat	tggcaccact	ctgtgtccct	ggaatgctag	ttggagtta	1740
aaatctcttg	aacagatttg	gaataacatg	acctggatgg	agtgggacag	agaaattaac	1800
aattacacaa	gcttaataca	ctccttaatt	gaagaatcgc	aaaaccagca	agaaaagaat	1860
gaacaagaat	tatttgaatt	agataaatgg	gcaagtttgt	ggaatttggt	taacataaca	1920
aatttggctgt	ggtatataaa	attattcata	atgatagtat	gaggcttggt	aggttttaaga	1980
atagtttttg	ctgtactttc	tatatgtaat	agagttaggc	agggatattc	accattatcg	2040
tttcagaccc	acctcccaat	cccgagggga	ccgcacaggc	ccgaaggga	agaagaagaa	2100

2148

<220>
<223> Description of the artificial sequence: synthetic DNA

```
<220>  
<221> misc_feature  
<222> (1377)..(1379)  
<223> env AGT, gp120 start
```

```
<220>
<221> misc_feature
<222> (1397)..(1403)
<223> BstEII cleavage site
```

```
<220>
<221> misc_feature
<222> (3537)..(3542)
<223> BamHI cleavage site
```

```
<220>
<221> misc_feature
<222> (3855)..(3857)
<223> env TAA, stop
```

<400>	10					
ctgacgcgcc	ctgtagcgggc	gcattaagcgc	cggcggggtgt	ggtgggttacg	cgcagcgtga	60
ccgctacact	tgccagcggc	ctagcgcgcc	ctccttttcgc	tttcttccc	tcctttctcg	120
ccagcttcgc	cggctttccc	cgtcaagctc	taaatcgggg	gctccccitta	gggtttccgat	180
ttagtctctt	acggcacctc	gaccocaaaa	aacttgattga	gggtgatagg	tcacgtatgt	240
ggccatcgcc	ctgatagacg	gttttttcgcc	ctttgacgtt	ggagtccacg	ttctttaata	300
gtggactctt	gttccaaact	ggaacaacac	tcaaccttat	ctcgggtctat	tccttttgatt	360
tataagggat	ttttccgatt	tcggcctatt	gggtaaaaaa	tgagctgatt	taacaaaaat	420
ttaacgcgaa	tttttaacaaa	atattaacgc	ttacaatttc	cattcgccat	tcaggctgcg	480
caactgttgg	gaaggggcgat	cgggtgcgggc	ctcttcgcta	ttacgccagc	tggcgaaagg	540
gggatgtgct	gcaaggcgat	taagttgggt	aacgccaggg	ttttcccagt	cacgacgttg	600
taaaacgcag	gccagtgagc	gtctagttat	taatagtaat	caattacggg	gtcattagtt	660
catagcccat	atatggagtt	ccgcgttata	taacttacgg	taaatggccc	gcctggctga	720
ccgcccacac	accccggccc	attgacgtca	ataatgcagt	atgttcccat	agtaacgcca	780
atagggactt	tccattgacg	tcaatgggtg	gagtatattac	ggtaaactgc	ccactttggca	840
gtacatcaag	tgtatcatat	gccaaagtac	ccccctattg	acgtcaatga	cggtaaatgg	900
cccgcctggc	attatgccca	gtacatgacc	ttatgggact	ttcttacttg	gcagtatcat	960
tcagtattag	tcatcgctat	taccattggg	atgcggtttt	ggcagtagat	caatgggctg	1020
ggatagcgg	ttgactcacg	gggattttcca	agtctccacc	ccattgacgt	caatggggagt	1080
ttgtttttgg	accaaaaatca	acgggacttt	ccaaaatgtc	gtaacaactc	cgccttattg	1140
acgcaaatgg	gcggttaggcg	tgtacgggtg	gaggtctata	taagcagagc	tcgtttagtg	1200
aaccgtcaga	tcgcctggag	acgccatoca	cgcgtttttg	acctccatag	aagacaccgg	1260
gacaattcga	gctcgggtacc	gtcgcacgcca	ccatgagagt	gaaggagaag	tatcagcact	1320
tgtggagatg	gggggtggaaa	tggggcacca	tgctccttgg	gatatttgat	atctgtatgt	1380
ctacagaaaa	atttgtgggtc	accgtctatt	atgggggtacc	tgatgtggaag	gaagcaacca	1440
ccactctatt	ttgtgcatac	gatcgtaaa	catatgatac	agaggtagat	aatgttttgg	1500
ccacacatgc	ctgtgtaccc	acgagcccca	accacaaga	agtagtattg	gtaaatgtga	1560
cagaaaaatt	taacatgtgg	aaaaatgaca	tggtagaaca	gatgcctgag	gatataatca	1620
gtttatggga	tczaagccta	aagccatgtg	taaaattaac	cccactctgt	gttagtttaa	1680
agtgacactg	tttgaagaat	gatacataata	ccaatagtag	tagcgggaga	atgataattg	1740
agaaaggaga	gataaaaaaac	tgtcttttca	atatcagcac	aagcataaga	gataaggtgc	1800

agaaagaata	tgrattcttt	tataaacttg	atatagtacc	aatagataat	accagctata	1860
ggttgataag	ttgtaacacc	tcagtcatta	cacaggcctg	tccaaaggta	tcctttgagc	1920
caattcccat	acattattgt	gccccggctg	gttttgcat	tctaaaatgt	aataataaga	1980
cgttcaatgg	aacaggacca	tgtacaaat	tcagcacagt	acaatgtaca	catggaatca	2040
ggccagtagt	atcaactcaa	ctgctgttaa	atggcagctc	agcagaagaa	gatgtagtaa	2100
ttagatctgt	caatttcaca	gacaatgcta	aaaccataat	agtcacagctg	aacacatctg	2160
tagaaattaa	ttgtacaaga	cccaacaaca	atacaagaaa	aagtatccgt	atccagaggg	2220
gaccagggag	agcattttgt	acaattaggaa	aaataggaaa	tatgagacaa	gcacattgta	2280
acattagtag	agcaaaaatgg	aatgccattc	taaaacagat	agctagcaaa	ttaagagaac	2340
aatttggaaa	taataaaaaca	ataatcttta	agcaatcctc	aggaggggac	ccagaaaattg	2400
taacgcacag	ttttaattgt	ggaggggaat	ttttctactg	taattcaaca	caactgttta	2460
atagtacttg	gtttaatagt	acttggagta	ctgaagggtc	aaataacact	gaaggaagtg	2520
acacaatcac	actcccatgc	agaataaaaac	aattttataaa	catgtggcag	gaagtaggaa	2580
aagcaatgta	tgccccctcc	atcagttggac	aaatttagat	ttcatcaaat	attactgggc	2640
tgtctattaac	aagagatggg	ggtaataaaca	acaatgggctc	cgagatcttc	agacctggag	2700
gaggcgatat	gaggggacaat	tggagaagtg	aattatataa	atataaagta	gtaaaaatttg	2760
aaccattagg	agtagcaccc	accaaggcaa	agagaagagt	ggtgcagaga	gaaaaaagag	2820
cagtgggaat	aggagctttg	ttccttgggt	tcttggggagc	agcaggaagc	actatgggct	2880
gcacgtcaat	gacgtctggc	gtacaggcca	gacaattatt	gtctgataata	gtgcagcagc	2940
agaacaattt	tgtgagggct	attgaggcgc	aacagcatct	gttgcaactc	acagcttggg	3000
gcatcaaaca	gctccaggca	agaatcctgg	ctgtggaaag	atacctaaag	gatcaacagc	3060
tcctgggggat	ttgggggttg	tctggaaaac	tcatttgcac	cactgctgtg	ctttggaatg	3120
ctagtgggag	taataaatct	ctggaacaga	tttggaaata	catgacctgg	atggagtggg	3180
acagagaaat	taacaaattac	acaagcttaa	tacactcctt	aattgaagaa	tcgcaaaacc	3240
agcaagaaaa	gaatgaacaa	gaattattgg	aattagataa	atgggcgaagt	ttgttggaatt	3300
ggtttaacat	aacaaattgg	ctgtggtata	taaaaattat	cataatgata	gtaggaggct	3360
tggtagggtt	aagaatagtt	tttgctgtac	tttctatagt	gaatagagtt	aggcagggat	3420
attcaccatt	atcgcttcag	accacactcc	caatcccag	gggacccgac	aggcccgaag	3480
gaatagaaga	agaaggttga	gagagagaca	gagacagatc	cattcgatta	gtgaacggtat	3540
ccttagcact	tatctgggac	gatctcgga	gcctgtgcct	cttcagctac	caccgcttga	3600
gagacttact	cttgatttga	acgaggattg	tggaaactct	gggacgcagg	gggtgggaag	3660
ccctcaaata	ttggtgggaat	ctcctacagt	attggagtca	ggaactaaag	aatagtgctg	3720
ttaaacttgc	caatgccaca	gccatagcag	tagctgaggg	gacagatagg	gttatagaag	3780
tattacaagc	agcttataga	gctattcgcc	acatacctag	agaataaaga	cagggcttgg	3840
aaaggatttt	gctataagat	gggtggcga	tggtaaaaa	gtagtgtgat	tggatggcct	3900
gctgtaagg	aaagaatgag	acgagctgag	ccagcagcag	atgggggtgg	agcagtatct	3960
cgagatctag	actagaacta	gcttcgatcc	agacatgata	agatacattg	atgagtttgg	4020
acaaaccaca	actagaatgc	taagtaaaaa	atgcttttat	tgtgaaattt	gtgatgctat	4080
ctctttat	gtaaccatta	taagctgcaa	taaacaaagt	aaacaaca	attgcattca	4140
ttttatgttt	cagggttcagg	gggaggtgtg	ggaggttttt	taaagcaagt	aaaacctcta	4200
caaatgtgtg	atggctgatt	atgattcctg	ctcgcgcggt	tcgggtgatga	cggtgaaaac	4260
ctctgacaca	tgragctccc	ggagacggtc	acagcttgtc	tgtaaagcga	tgccggggagc	4320
agacaagccc	gtcagggcgc	gtcagcgggt	gttggcgggt	gtcggggcgc	agccatgacc	4380
cagtcacgta	gcgatagcgg	agtgtatact	ggcttaacta	tgcggcatca	gagcagattg	4440
tactgagagt	gcaccatatt	tcgggcccgc	ttgctggcgt	ttttccatag	gtccgcctcc	4500
cctgacgagc	atcacaaaaa	tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	4560
taaagatacc	aggcgtttcc	ccctgggaag	tcctctgtgc	gctctcctgt	tcggaccttg	4620
ccgcttaccg	gatacctgtc	cgcctttctc	ccctcgggaa	cgggtggcgt	ttctcatagc	4680
tcacgctgta	ggtatctcag	ttcgggtgtg	gtcgttctgt	ccaagctggg	ctgtgtgcac	4740
gaaccccccg	ttcagcccca	ccgctcgccc	ttatccggta	actatcgtct	tgagccaac	4800
ccggtaagac	acgacttatc	ggcactggca	gcagccactg	gtaacaggat	tagcagagcg	4860
aggtagtagc	gcggtgtctc	agagttcttg	aagtcgttgc	ctaactacgg	ctacactaga	4920
aggacagtat	ttggttatctg	cgtctgtgtg	aagccagtta	ccctcggaaa	aagagtttgt	4980
agctcttgat	ccggcaaaaa	aaccaccgct	ggtagcgggt	gtttttttgt	ttgcaagcag	5040
cagattacgc	gcagaaaaaa	aggatctcaa	gaagatcctt	tgatcttttc	tacgggggtct	5100
gacgtctcag	ggaacgaaaa	ctcacgttaa	gggatttttg	tcatgagatt	atcaaaaaag	5160
atcttcacct	agatcctttt	aaatataaaa	tgaagtttta	aatcaatcta	aagtatatat	5220
gagtaaacct	ggtctgacag	ttaccaatgc	ttaatcagtg</			


```
tgtatgctggc gaccgagttg ctcttgcccg gcgtcaatac gggataatac cgcgccacat 5880
agcagaactt taaaagtgt catcattgga aaacgttctt cggggcgaaa actctcaagg 5940
atcttaccgc ttttgagatc cagttcgatg taaccactc gtgcacccaa ctgatcttca 6000
gcatctttta ctttcaccag cgtttctggg tgagcaaaaa caggaaggca aaatgccgca 6060
aaaaagggaa taaggcgac acggaaatgt tgaatactca tactcttctt ttttcaatat 6120
tattgaagca tttatcaggg ttattgtctc atgagcggat acatatttga atgtatttag 6180
aaaaataaac aaataggggg tccgcgcaca tttccccgaa aagtggccac 6229
```

```
<210> 11
<211> 860
<212> DNA
<213> Human immunodeficiency virus

<220>
<221> misc_feature
<222> (1)..(860)
<223> PI-932 original sequence V1-V2-V3-loop
```

```
<400> 11
tgtgtaccca cagaccccaa cccacaaaag gtagtattgg aaaatgtgac agaaaatttt 60
aacatgtgga aaatgacat ggtagaacag atgcatgagg atataatcaa tttatgggat 120
caaagcctta agccatgtgt aaaactaacc ccactctgtg ttacttttaa ttgactgat 180
gctgatttaa attgcaataa tactgattta aattgcacta aagctaattt ggggaaaaat 240
actcataaca atactattag tgggaaaata atagagaaag tagaaataaa aaactgctct 300
ttcaagggtca ccacaggcat aagggtataag atgcaaaaag aatatgcact tttgaataaa 360
cttgatatag taccaataga taatgataag aataatacta actttatatt gataagttgt 420
aacacctoga ccattacaca ggcctgtcca aaggtatcct ttgagccaat tcccatatat 480
ttttgtgccc cggctgggtt tgcgattcta aagtgtaatg aaaagagtta cagtggaaaa 540
ggaccatgta aaaatgtcag cacagtacaa tgtacacatg gaattaggcc agtagtgtca 600
actcaactgc tgttgaatgg cagtctagca gaaaaagaag tagtaattag atctgagaat 660
ttcacagaca atgctaaaac cataatagta cagctgaagg aatctgtaaa cattacttgt 720
ataagacccc acaacactgt aacagacagg atacatatag ggccaggagg atcatttcac 780
acaacaagaa aaataaaaagg agatataaga caagcacatt gtagccttag gagaaaagat 840
tggaataaca ctttacaaga gatagctagc 860
```

```
<210> 12
<211> 870
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> Description of the artificial sequence: PI-932
gene cassette, comprising the cleavage sites for
restriction enzymes BspTl, PstI, BclI, EcoRI,
BglII, PvuII, XbaII, NheI
```

```
<400> 12
tgtgtaccca cagaccccaa cccacaaaag gtagtattgg aaaatgtgac agaaaatttt 60
aacatgtgga aaatgacat ggtagaacag atgcatgagg atataatcaa tttatgggat 120
caaagcctta agccatgtgt aaaactaacc ccactctgtg ttacttttaa ttgactgat 180
gctgatttaa attgcaataa tactgattta aattgcacta aagctaattt ggggaaaaat 240
actcataact gcagtattag tgggaaaata atagagaaag tagaaataaa aaactgctct 300
ttcaagggtca ccacaggcat aagggtataag atgcaaaaag aatatgcact tttgaataaa 360
cttgatatag taccaataga taatgataag aataatacta actttatatt gataagttgt 420
aacacctcgg tgatcacaca ggcctgtcca aaggtatcct ttgagccaat tcccatatat 480
ttttgtgccc cggctgggtt tgcgattcta aagtgtaatg aaaagagtta cagtggaaaa 540
ggaccatgta aaaatgtcag cacagtacaa tgtacacatg gaattcggcc agtagtgtca 600
actcaactgc tgttgaatgg cagtctagca gaaaaagaag tagtaattag atctgagaat 660
ttcacagaca atgctaaaac cataatagta cagctgaagg aatctgtaaa cattacttgt 720
ataagacccc acaacactgt aacagacagg atacatatag ggccaggagg atcatttcac 780
acaacaagaa aaataaaaagg agatataaga caagcacatt gtagcctttc tagaaaagat 840
tggaataaca ctttacaaga gatagctagc 870
```

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of

SCHREIBER

Atty. Ref.: 35-213

Serial No. Unknown

Group:

U.S. National Phase of PCT/EP99/09759

Filed: August 10, 2001

Examiner:

For: VIRAL VACCINE

* * * * *

August 10, 2001

Assistant Commissioner for Patents
Washington, DC 20231

Sir:

PRELIMINARY AMENDMENT

In order to place the above-identified application in better condition for examination, please amend the application as follows:

IN THE CLAIMS

Please substitute the following amended claims for corresponding claims previously presented. A copy of the amended claims showing current revisions is attached.

3. (Amended) Protein vaccine according to claim 1 , characterized in that it comprises a mixture of GP120 proteins of HIV which in each case differ from each other in their amino acid sequence in the region of the V2 loop and/or of the V3 loop.

6. (Amended) DNA vaccine according to claim 4 , characterized in that the mixture contains $\geq 10^3$ and preferably $\geq 10^4$ DNA molecules which differ from each other in their nucleic acid sequence.

7. (Amended) DNA vaccine according to claim 4, characterized in that it codes for a mixture of structurally different GP120 proteins of HIV, in which the vaccine contains a mixture of DNA molecules, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop.

11. (Amended) Nucleic acid sequence according to claim 9 , characterized in that the sequence is modified by the introduction of silent mutations.

12. (Amended) Nucleic acid sequence according to claim 9, characterized in that it contains the sequence given in SEQ ID NO: 9.

17. (Amended) Nucleic acid sequence according to claim 15 or a single-stranded nucleic acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12 , characterized in that the fragment/the fragments contain(s) inosine, a nucleic acid exchange or a mutation at 9 to 20 positions.

18. (Amended) Double-stranded DNA which comprises hybrids of the single-stranded nucleic acid sequence according to claim 15 with a single-stranded nucleic

acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12.

27. (Amended) Expression vector, characterized in that it contains an inserted nucleic acid sequence according to claim 9.

33. (Amended) Vector mixture according to claim 30, characterized in that the plasmids can be expressed in *E. coli* as host cell.

34. (Amended) Vector mixture according to claim 30, characterized in that the plasmids can be expressed in eukaryotic cells, preferably in Cos, CHO or BHK cells, as host cells.

40. (Amended) Process according to claim 38, characterized in that the nucleic acid sequence coding for a viral protein is the sequence according SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same.

41. (Amended) Process for the preparation of the vector mixture according to claim 33, characterized in that plasmids, the nucleic acid sequences of which in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop in each case through random distribution of the bases at the

varied nucleotide positions, are ligated into a vector which can be expressed in host cells.

43. (Amended) Process for the preparation of the host cells composing transforming *E.coli*. with a vector mixture according to claim 30 .

44. (Amended) Process for the preparation of a protein vaccine which comprises a mixture of viral protein molecules, characterized in that the molecules are sequence variants of a single viral protein or of part of same, the mixture containing $\geq 10^2$ sequence variants, which is obtainable by expression of a plasmid-DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations , said process comprising cultivating host cells according to claim 35 under conditions which allow the expression of the mixture of viral protein sequence variants.

45. (Amended) Process for the preparation of a DNA vaccine which codes for a mixture of structurally different virus proteins, characterized in that the vaccine contains a mixture of sequence variants of a viral DNA molecule or of part of same, the mixture containing $\geq 10^2$ DNA molecules which differ from each other in their nucleic acid sequence, where the mixture, because of the variation of nucleotide positions, contains randomly distributed sequence combinations wherein said process is carried out according to claim 41 , wherein the plasmids are ligated into a vector which can be expressed in host cells of the organism to be vaccinated.

46. (Amended) A method of preparing a vaccine comprising forming a mixture of structurally different viral proteins which are sequence variants of a viral protein or of part of same, for the prevention and/or therapy of a virus infection in humans.

47. (Amended) A method of preparing a vaccine comprising forming a mixture according to claim 23 for the prevention and/or therapy of a HIV infection in humans.

48. (Amended) A method of preparing a vaccine comprising forming a mixture of DNA molecules which code for sequence variants of a viral protein or of part of same, for the prevention and/or therapy of a virus infection in humans.

49. (Amended) A method of preparing a vaccine comprising forming a nucleic acid mixture according to claim 19 for the prevention and/or therapy of a virus infection in humans.

50. (Amended) A method of preparing a vaccine comprising forming the nucleic acid mixture according to claim 19 for the expression in host cells selected from the group consisting of *E. coli*, Cos, CHO and BHK cells.

51. (Amended) A method of producing a protein mixture which comprises sequence variants of the GP120 protein, characterized in that it is a mixture of proteins which contain amino acid sequences which in each case differ from each other in the V2 loop and/or in the V3 loop, the mixture being obtainable by expression of a plasmid DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations; said method comprising expressing a vector mixture which contains a mixture of plasmids which contains an inserted double-stranded DNA which comprises hybrids of the single-stranded nucleic acid sequence according to claim 15 with a single-stranded nucleic acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12; characterized in that the nucleic acid sequences of the plasmids differ in each case from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop, where the mixture of the plasmids, because

of the variation of nucleotide positions, contains randomly distributed sequence combinations .

52. (Amended) A method of preparing a protein mixture which comprises sequence variants of the GP120 protein, characterized in that it is a mixture of proteins which contain amino acid sequences which in each case differ from each other in the V2 loop and/or in the V3 loop, the mixture being obtainable by expression of a plasmid DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations, said method comprising culturing a host cell according to claim 35 .

54. (Amended) Pharmaceutical composition for the prevention and/or therapy of a virus infection, characterized in that it comprises a protein mixture and a nucleic acid mixture, the protein mixture comprising sequence variants of a viral protein or of part of same, and the nucleic acid mixture comprising DNA molecules which code for sequence variants of a viral protein or of part of same, said composition comprising a protein mixture according to claim 23 and a nucleic acid mixture which comprises double-stranded DNAs, the nucleic acid sequences of which are derived from the *env* sequence in SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same, characterized in that the nucleic acid sequences in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop .

REMARKS

The above amendments are made to place the claims in a more traditional format. Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached pages are captioned "**Version With Markings To Show Changes Made.**"

SCHREIBER

Serial No. **Unknown**

U.S. National Phase of PCT/EP99/09759

Respectfully submitted,

NIXON & VANDERHYE P.C.

By: _____



B. J. Sadoff
Reg. No. 36,663

BJS:ms

1100 North Glebe Road, 8th Floor

Arlington, VA 22201-4714

Telephone: (703) 816-4000

Facsimile: (703) 816-4100

VERSION WITH MARKINGS TO SHOW CHANGES MADE

IN THE CLAIMS

3. (Amended) Protein vaccine according to claim 1 [or 2], characterized in that it comprises a mixture of GP120 proteins of HIV which in each case differ from each other in their amino acid sequence in the region of the V2 loop and/or of the V3 loop.

6. (Amended) DNA vaccine according to claim 4 [or 5], characterized in that the mixture contains $\geq 10^3$ and preferably $\geq 10^4$ DNA molecules which differ from each other in their nucleic acid sequence.

7. (Amended) DNA vaccine according to [one of claims 4 to 6] claim 4, characterized in that it codes for a mixture of structurally different GP120 proteins of HIV, in which the vaccine contains a mixture of DNA molecules, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop.

11. (Amended) Nucleic acid sequence according to claim 9 [or 10], characterized in that the sequence is modified by the introduction of silent mutations.

12. (Amended) Nucleic acid sequence according to [claims 9 to 11] claim 9, characterized in that it contains the sequence given in SEQ ID NO: 9.

17. (Amended) Nucleic acid sequence according to claim 15 or [16] a single-stranded nucleic acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-

long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12, characterized in that the fragment/the fragments contain(s) inosine, a nucleic acid exchange or a mutation at 9 to 20 positions.

18. (Amended) Double-stranded DNA which comprises hybrids of the single-stranded nucleic acid sequence according to claim 15 [or 17] with [the] a single-stranded nucleic acid sequence [according to claim 16 or 17] which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12.

27. (Amended) Expression vector, characterized in that it contains an inserted nucleic acid sequence according to [claims 9 to 14]claim 9.

33. (Amended) Vector mixture according to [one of claims 30 to 32]claim 30, characterized in that the plasmids can be expressed in *E. coli* as host cell.

34. (Amended) Vector mixture according to [one of claims 30 to 32]claim 30, characterized in that the plasmids can be expressed in eukaryotic cells, preferably in Cos, CHO or BHK cells, as host cells.

40. (Amended) Process according to claim 38 [or 39], characterized in that the nucleic acid sequence coding for a viral protein is the sequence according SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same.

41. (Amended) Process for the preparation of the vector mixture according to [claims 33 and 34]claim 33, characterized in that plasmids, the nucleic acid sequences of which in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop in each case through random distribution of the bases at the varied nucleotide positions, are ligated into a vector which can be expressed in host cells.

43. (Amended) Process for the preparation of the host cells [according to claim 35 or 36, characterized in that the host cells are transformed] composing transforming *E.coli*, with a vector mixture according to [claims] claim 30 [to 32].

44. (Amended) Process for the preparation of a protein vaccine which comprises a mixture of viral protein molecules, characterized in that the molecules are sequence variants of a single viral protein or of part of same, the mixture containing $\geq 10^2$ sequence variants, which is obtainable by expression of a plasmid-DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations [according to one of claims 1 to 3, characterized in that the host cells are cultivated] , said process comprising cultivating host cells according to [one of claims 35 to 37] claim 35 under conditions which allow the expression of the mixture of viral protein sequence variants.

45. (Amended) Process for the preparation of a DNA vaccine which codes for a mixture of structurally different virus proteins, characterized in that the vaccine contains a mixture of sequence variants of a viral DNA molecule or of part of same, the mixture containing $\geq 10^2$ DNA molecules which differ from each other in their nucleic acid sequence, where the mixture, because of the variation of nucleotide positions, contains

randomly distributed sequence combinations wherein said [according to [one of claims 4 to 8, characterized in that the] process is carried out according to claim 41 [or 42],
wherein the plasmids [according to the invention being] are ligated into a vector which can be expressed in host cells of the organism to be vaccinated.

46. (Amended) A method of preparing a vaccine comprising forming a [Use of a] mixture of structurally different viral proteins which are sequence variants of a viral protein or of part of same, for [the preparation of a vaccine for] the prevention and/or therapy of a virus infection in humans.

47. (Amended) A method of preparing a vaccine comprising forming a mixture [Use of a protein mixture] according to [one of claims 23 to 25]claim 23 [for the preparation of a vaccine] for the prevention and/or therapy of a HIV infection in humans.

48. (Amended) A method of preparing a vaccine comprising forming [Use of] a mixture of DNA molecules which code for sequence variants of a viral protein or of part of same, for [the preparation of a vaccine for] the prevention and/or therapy of a virus infection in humans.

49. (Amended) A method of preparing a vaccine comprising [Use of] forming a nucleic acid mixture according to [claims 19 to 22]claim 19 for the [preparation of a vaccine for the] prevention and/or therapy of a virus infection in humans.

50. (Amended) A method of preparing a vaccine comprising [Use of] forming the nucleic acid mixture according to [one of claims 19 to 22]claim 19 for the [preparation of a vector mixture according to one of claims 30 to 32 which can be expressed]
expression in host cells[, the host cells being] selected from the group consisting of *E. coli*, Cos, CHO and BHK cells.

51. (Amended) A method of producing a protein mixture which comprises
sequence variants of the GP120 protein, characterized in that it is a mixture of proteins

which contain amino acid sequences which in each case differ from each other in the V2 loop and/or in the V3 loop, the mixture being obtainable by expression of a plasmid DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations; said method comprising [Use of the vector mixture according to one of claims 30 to 32] expressing a vector mixture which contains a mixture of plasmids which contains an inserted double-stranded DNA which comprises hybrids of the single-stranded nucleic acid sequence according to claim 15 with a single-stranded nucleic acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, characterized in that in the V3 loop a 247 bp-long Bg1II-XbaI fragment or a 283 bp-long Bg1II-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-Bc1I fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to a nucleic acid sequence characterized in that it contains the sequence given in SEQ ID NO: 12; characterized in that the nucleic acid sequences of the plasmids differ in each case from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop, where the mixture of the plasmids, because of the variation of nucleotide positions, contains randomly distributed sequence combinations [for the expression of a protein mixture according to one of claims 23 to 25].

52. (Amended) A method of preparing a protein mixture which comprises sequence variants of the GP120 protein, characterized in that it is a mixture of proteins which contain amino acid sequences which in each case differ from each other in the V2 loop and/or in the V3 loop, the mixture being obtainable by expression of a plasmid DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations, said method comprising culturing a [Use of the] host cell according to [one of claims 35 to 37]claim 35 [for the preparation of a protein mixture according to one of claims 23 to 25].

SCHREIBERSerial No. **Unknown**

U.S. National Phase of PCT/EP99/09759

54. (Amended) Pharmaceutical composition for the prevention and/or therapy of a virus infection, characterized in that it comprises a protein mixture and a nucleic acid mixture, the protein mixture comprising sequence variants of a viral protein or of part of same, and the nucleic acid mixture comprising DNA molecules which code for sequence variants of a viral protein or of part of same, said composition comprising [according to claim 53, characterized in that it comprises] a protein mixture according to claim 23 [claims 23 to 25] and a nucleic acid mixture which comprises double-stranded DNAs, the nucleic acid sequences of which are derived from the *env* sequence in SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same, characterized in that the nucleic acid sequences in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop [according to claims 19 to 22].

09/913159

518 Rec'd PCT/PTO 10 AUG 2001

Strathmann AG & Co.

Sellhopsweg 1

22459 Hamburg

(P 52348 WE/mo/wo)

01.12.1999

VIRUS VACCINE

09/913159

The present invention relates to a pharmaceutical composition or a vaccine which comprises a mixture of viral protein molecules which are sequence variants of a single viral protein or of part of same. The invention furthermore relates inter alia to a DNA vaccine which codes for a mixture of structurally different virus proteins, the vaccine containing a mixture of sequence variants of a viral DNA molecule or of part of same, which code for sequence variants of a viral protein or part. According to a preferred version of the invention, the viral proteins are sequence variants of the GP120 protein of the human immunodeficiency virus (HIV) which differ from each other in their amino acid sequence in the region of the V2 loop and/or of the V3 loop, preferably both the V2 and the V3 loop. The invention furthermore relates to the preparation of the virus vaccines including the intermediate stages or constructs, preparation processes and uses connected with them.

In many viral infections, in particular in the HIV-1 infection, a strong immune defence is observed, which is in

a position to control the virus over a period of several years. The period in which the virus is controlled and no illness symptoms are observed, is called the asymptomatic phase of the (HIV) illness. During the course of the illness, new virus variants repeatedly form. It is thereby possible for the virus to escape the human immune defence and to repeatedly infect new defence cells of the immune system (cf. M. Schreiber et al., J. Virol. 68 No. 6 (1994) 3908-3916; J. Gen. Virol. 77 (1996) 2403-2414; Clin. Exp. Immunol. 107 (1997) 15-20; J. Virol. 71 No. 12 (1997) 9198-9205).

In the prior art, for the treatment of viral illnesses, such as e.g. the polio virus (Heraud F et al., Biologicals, 1993, 21:311-316), Hanta virus (Ulrich R et al., 1998 Vaccine 16:272-280; Schmaljohn CS et al., 1992 vaccine 10:10-13), Lassa virus (Morrison HG et al., 1989 Virology 171:179-188; Clegg JC et al., 1987 Lancet 2:186-188), hepatitis A virus (Clemens et al., 1995 J. Infect. Dis. 171:Suppl.1:p.44-p.49; Andre et al., 1990 Prog. Med. Virol. 37:72-95) and hepatitis B virus infection (McAleer et al., 1984 Nature 307:178-180) but also the HIV infection (Egan et al., 1995 J. Infect. Dis. 171:1623-1627, Kovac et al., 1993 J. Clin. Invest. 92:919-928), only single, genetically unmodified, specific virus antigens or single inactivated virus strains have been used for studies of suitable vaccination strategies. In the case of HIV, for example, the external envelope proteins of two virus strains were up until now prepared for vaccine experiments on humans (MN and SF2) (Zolla-Pazner et al., J. Infect. Dis. 178 (1998) 1502-1506). Both GP120 molecules differ from each other in the amino acid sequence, but particularly in the amino acid

sequence of the variable regions, such as e.g. of the V3 loop. Because of the different V3 loop sequences, the two virus strains have different phenotypic properties. The HIV-1 MN strain is a virus variant which preferentially infects macrophages and cells which possess the viral co-receptor CCR5. Viruses of the SF2 type, on the other hand, multiply preferentially in T cells and use the viral co-receptor CXCR4. Such viruses are therefore also called T cell trophs (e.g. HIV strain) or macrophagotrophs (e.g. HIV strain MN). Vaccination experiments have been carried out in chimpanzees with these two GP120 variants. It has been shown in these experiments that an immune response can be induced which not only protects against the two HIV strains MN and SF2, but is also capable of preventing infection by other virus strains, but also by HIV-1 patient isolates. In contrast to chimpanzees, only one of the two GP120 variants has been used in humans for vaccination experiments up until now. Neither MN GP120 nor SF2 GP120 offers certain protection against infection by a virus variant such as occurs in a person infected by HIV-1 (patient isolate or wild-type virus). J. A. Levy offers in *"HIV and the Pathogenesis of AIDS"*, Publisher: Jay. A. Levy, Chapter 15, 2nd Edition, ASM Press Washington, D.C., 1998, an overview of the state of research in connection with vaccination with GP120 envelope protein.

The disadvantage of the vaccination strategies discussed or examined up until now in the prior art is, among other things, that, with the vaccines used, one is not in a position to prevent the formation of ever-new virus variants during the course of the viral illness.

The object of the present invention is therefore to provide a vaccine which in particular is in a position to prevent or clearly reduce the formation of new virus variants during the course of a viral illness and to limit or prevent the possibilities that the virus can escape the human immune defence and repeatedly infect new defence cells of the immune system.

In order to achieve the object, the items described in the following claims are proposed according to the invention.

The subject of the present invention is thus inter alia a protein vaccine which comprises a mixture of viral protein molecules, the molecules being sequence variants of a single viral protein or of part of same.

According to the invention, sequence variants of a protein are understood to be those molecules which contain an amino acid sequence derived vis-à-vis a native viral protein or a part (fragment) of same, the variants differing from each other in that at least one amino acid can be exchanged at any chosen point of the sequence or parts of same. The sequence variants preferably contain several amino acid exchanges at various points of the sequence, which are responsible for the production but also the bonding of virus-neutralizing antibodies. The number and position of the exchanged amino acids depends on the amino acid variability of the regions of the gp120 which are observed during the cell culture of adapted and wild-type HIV isolates. According to the invention, the sequence variants have a heterogeneity at at least two amino acid positions of the sequence or parts of same. A heterogeneity at three

100120" 65121500

to eight and preferably at more than eight amino acid positions is particularly preferred, all amino acids occurring at these positions being possible. The number of possible sequence variants which represent the vaccine in their totality result from the combination of the various amino acids which are possible per position.

The invention relates in particular to a vaccine which comprises a mixture of $\geq 10^2$ sequence variants, i.e. a mixture of more than 10^2 molecules of various amino acid sequences (homologues). A vaccine is particularly preferred which comprises $\geq 10^3$ and preferably $\geq 10^4$ sequence variants.

Within the framework of the present invention it is assumed that the vaccine can comprise, along with the sequence variants, the protein per se from which the sequence variants are derived.

An anti-viral active ingredient is thus provided according to the invention which comprises virus-specific proteins which all differ in their sequences or parts of same. In order to achieve this, the protein-coding gene is newly synthesized in order to generate new cleavage sites for DNA-restriction enzymes, in order to permit the exchange of specific regions. By chemical synthesis of DNA fragments which code for the concerned protein section, a gene bank is then produced for the protein-coding sequence. The expression vectors which code for the protein, are then transfected as a mixture into cells. From these protein-producing cells the mixture of the various viral proteins

can then be isolated which according to the invention represents the preferred active ingredient.

In the case of HIV as a viral illness, the chronic infection and the associated continuous damaging of the immune system during the course of the illness lead to the complete loss of the specific virus defence. Neutralizing antibodies which have the property of forming a specific, very strong bond with certain antigenic structures are part of the specific virus defence. Foreign antigens are thereby marked and their interaction with e.g. virus receptors are blocked. Such a specific virus defence are neutralizing antibodies against the V3 loop of the external GP120 envelope protein. Such anti-V3 loop antibodies are in a position to prevent the infection of cells. In animal models, it has been shown that an infection by HIV-1 can be prevented by administering a specific monoclonal V3 loop antibody. By administering the same monoclonal antibody, it was also possible to heal an existing infection.

In contrast to experimental systems, the development of ever-new virus variants is however observed in the natural course of HIV infection. Thus hundreds of different V3 loop variants are found in a single patient at one time, as the variation of the HIV-1 is particularly high precisely in the V3 loop. The V3 loop is an important dominant antigenic domain of the GP120. A very specific humoral immune response is therefore formed against each V3 loop. The result of the induction of a highly specific immune response against the V3 loop is that neutralizing antibodies against the V3 loop of the HIV-1 variant A are not in a position to neutralize the variant B and vice

versa (Schreiber et al., J. Virol. 68 (1994) 3908-16, Abrahamsson et al., 4 (1990) 107-12).

In the asymptomatic phase of the HIV illness, the cell-free virus variants in the serum are neutralized by antibodies. Only later can virus variants be observed in the serum, which escape autologous neutralization. These V3 loop escape variants are no longer recognized by V3 loop antibodies and are thus no longer neutralized either. All other virus variants which are found in the serum of the patients are however recognized by autologous V3 loop antibodies. This indicates that during the course of the illness, virus variants occur against which no V3 loop antibodies exist. If the antibody content of serum samples of a patient is examined over a period of several years, the V3 loop antibodies against the V3 loop escape variant which occurs in the later stage can be detected in the serum samples which were taken at the beginning of the asymptomatic phase. The result is not, as with other infectious diseases, the standard escape of the pathogen through ever-new antigenic variation, but the switching-off of the neutralizing immune response to a virus replicating in the patient. The observed absence of the neutralizing V3 loop antibodies is thus the result of a continuous loss of the type-specific V3 loop antibodies. The corresponding virus can multiply through the loss of the neutralizing antibodies, which leads to an increase in the viral load in the serum of the patients. In the course of the illness, the increase in the viral load in the lymph nodes is also observed (Chun et al., Nature 387 (1997) 183-188).

A virus which establishes itself against the immune system in the patient and develops into the dominant virus variant must necessarily overcome all antiviral barriers which the immune system offers. Along with neutralizing antibodies, cytotoxic T cells are also in a position to suppress the virus reproduction. Cytotoxic T cells effect the death of HIV infected CD4+ T cells. Along with the loss of the neutralizing antibodies, the loss of such cytotoxic T cells against HIV infected cells is also observed (Zerhoui et al., Thymus 24 (1997) 203-219; Gould et al., Semin Cell Dev Biol 9 (1998) 321-328; Wagner et al., J Gen Virol 74 (1993) 1261-1269; O'Toole et al., AIDS Res Hum Retroviruses 8 (1992) 1361-1368; Shearer et al., 137 (1986) 2514-2521). The object of a heterologous vaccine is therefore to induce or activate the mixture of many different GP120 envelope proteins of HIV both neutralizing antibodies and cytotoxic T cells against many different virus variants.

As already stated, the HIV illness is characterized in that the virus is constantly changing. This is effected by the error rate of the viral enzyme reverse transcriptase which commits errors during the increase of the viral genetic information. Mutants are thereby produced which are on the one hand selected because of their biological properties and the antiviral action of the human immune system. As with HIV, there are also other pathogens which transcribe their genetic information without error correction, which can lead to the formation of many variants. These include Hanta and Lassa viruses as well as hepatitis A, B and C viruses. Thus it is observed in people vaccinated for hepatitis B that vaccination fails in approx. 2%. In these

2%, hepatitis B "escape" variants are found which cannot be suppressed by the immune response induced by the vaccine.

The formation of vaccine, therapy and immune escape variants is based on the genetic variability of these viruses (Blum, Int J Clin Lab Res 27 (1997) 213-214; Jongerius et al., Transfusion 38 (1998) 56-59).

Up until now it has not been known in the state of the art how such a loss of virus-neutralizing antibodies can be countered. The strategies adopted to date were successful only in certain respects, i.e. in relation to the formation and the maintenance of individual variant-specific antibodies against specific viruses, but the treatment methods proposed to date in the form of antiviral medicaments or inoculants based on a memo-substance cannot counter the loss of type-specific antibodies against the variety of the protein variants formed by the virus in the course of the illness.

It is possible, through the present invention of a heterogeneous vaccine, to prevent the loss of neutralizing V3 antibodies or at least to clearly counter this loss.

On the basis of the present invention, for the first time an immunoreconstitutive treatment of virus-infected patients, in particular infected with HIV-1, is provided in which the immune system is activated such that the naturally acquired immune defence against the virus population of the various virus variants is regenerated and stimulated anew, in order to thus prevent the virus variants present in the patient from multiplying.

In short, the invention is based on the concept of preparing a mixture of GP120 proteins for an active ingredient against HIV-1, these GP120 proteins differing e.g. either in the V2 or the V3 loop or simultaneously in the two variable domains, i.e. the V2 loop and the V3 loop. In order to achieve this, the GP120-coding gene has to be newly synthesized accompanied by the formation of new (monovalent) cleavage sites for DNA-restriction enzymes which allow the specific exchange of the e.g. V2 and V3 region. By chemical synthesis of DNA fragments which code for the V2 loop and V3 loop a V2/V3 gene bank for GP120 is then produced. The GP120 expressions vectors are finally transfected into cells as a mixture, and the mixture of the various GP120 proteins which can be used as an active ingredient for the prophylaxis and/or therapy of the HIV illness or AIDS can then be isolated from these GP120-producing cells.

The consideration on which the present invention is based thus comprises the preparation for an immunotherapy or a vaccine against HIV-1 of many different GP120 molecules which carry V3 loop sequences (and/or V2 loop sequences) such as can also be identified in virus variants in patients.

The production of a mixture of natural virus variants in the cell culture system is very expensive. It must be borne in mind in particular that the composition of a virus mixture changes as cells which can be infected by HIV-1 have to be used in the cell culture. Some virus variants which have selective advantages, e.g. faster growth

kinesis, therefore become dominant in the cell culture system after only a few days and expel the other slower virus variants. This applies above all when the various virus variants which are present in the peripheral mononuclear blood cells (PMBC) or in the serum of patients are to be isolated. As it is not possible to cultivate a constant mixture of HIV variants, neither can a corresponding mixture of GP120 proteins be produced nor isolated in this way from virus cultures.

Within the framework of the present invention, a genetic-engineering approach was therefore chosen for the preparation of HIV-1 variants and GP120 variants. For the preparation of virus variants and recombinant GP120, a cloning system was constructed which consists of two vectors. The central component of both vectors is a novel, chemically synthesized HIV-1 envelope protein gene (HIV-1 env gene) which codes for the GP120 protein. One vector contains the entire genome of HIV-1. After transfection of this vector into cells, the cells produce infectious virus particles. This facilitates the production of a mixture of virus variants as cells can be used which are resistant to an HIV-1 infection. Therefore, in such a cell culture system, the composition of the virus mixture cannot be changed, because of the biological properties of the virus variants. The second vector facilitates the expression of GP120-variants in eukaryotic cells. This vector serves directly for the preparation of the inoculant (the vaccine).

For the genetic-engineering preparation and expression of the V3 loop GP120 variants in eukaryotic cells, a special

gene construct for HIV-1-env was created, called gene cassette in the following. For the preparation of the gene cassette, according to the invention the entire coding sequence of the env gene was chemically synthesized. With the help of this method it is possible to change the coding sequence of the gene as desired, new DNA-recognition sequences for DNA-cutting restriction enzymes being able to be incorporated into the env-gene sequence. It must be borne in mind that the amino acid sequence of the original GP120 (preferably of the strains NL4-3 and PI-932) does not change, while new restriction cleavage sites form in the DNA sequence of the env gene. At the same time, according to the invention cleavage sites for enzymes which repeatedly occur in the env sequence are removed so that in each case there is only one cleavage site for a specific restriction enzyme, such as e.g. BglIII. According to a preferred version of the invention ten new, once-occurring (monovalent) cleavage sites are inserted at intervals of approx. 150 base pairs. The new env gene produced in this way is also called gene cassette within the framework of the present invention. In principle, the new env fragment is similar to a polylinker.

The cleavage sites for the restriction enzymes BstEII and BamHI define the gene cassette preferred according to the invention (SEQ ID NO: 9). In order to be able to exchange the region of the env gene, both cleavage sites are present only once in the pBSCenvATG expression vector (SEQ ID NO: 10) for gp120 and in retroviral vector pNL4-3. The vector, given the reference code pBSCenv-V3 by the depositor, was deposited on 6th January 1999 at the DSMZ - Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH,

Mascheroder Weg 1b, 38124 Brunswick, Germany, under the accession number DSM 12612 in accordance with the Budapest Treaty.

Through the exchange of the gene section defined by BstEII and BamHI, the newly created sections of the env gene which are likewise defined by BstEII and BamHI, can be cloned into both vectors. The PstI/BclI V2 loop and the BglII/XbaI V3 loop env fragment are chemically or enzymatically synthesized and then cloned into a standard vector such as e.g. pUC 18 or 19. In this vector, the fragment is checked by sequencing and then cut out of the vector by a Pst/BcII or BgIII/XbaI digestion and transferred into the pBSCenvATG and then into the pNL4-3 vector. In place of the intermediate cloning into a standard vector (pUC 18, pUC 19 etc.), the PstI/BclI V2 loop and BglII/XbaI V3 loop fragments can also be cloned directly into pBSCenvATG.

As there is a cleavage site for a restriction enzyme every ca. 100 base pairs in the gene cassette, all regions of the env gene, in particular the V3 loop or V2 loop, can be exchanged for any DNA fragments. The section coding for the V3 loop can be exchanged for a BglII-XbaI fragment, which has a size of 244 base pairs (cf. SEQ ID NO: 9; nucleotides 708 to 955). The aim is to exchange the gene sections which code for the variable loops of the GP120 protein for new DNA fragments. The fragments are synthetically produced and then cloned into the env gene cassette. If the sequence is varied at certain positions during the synthesis of the V2 loop and V3 loop DNA fragments, i.e. all four nucleotides are incorporated equally or the nucleotide inosine is used at this position, env-gene variants with a pre-set

variation of the thereby-coded amino acid sequence can be produced. The V3 loop sequences of patient isolates are the starting point for the preparation of the GP120 mixture. If these variants are cloned into the *gp120* gene cassette, the corresponding GP120 proteins can be expressed with the variable antigenic domains in eukaryotic cells. Sequence data of HIV-1 patient isolates are available for the variation of the V3 loop region of the *env* gene. Patient isolates are virus variants which have been cultivated direct from patient material, serum or infected cells from the blood in the laboratory. These viruses are characterized, unlike HIV-1 viruses which are cultivated in the laboratory for a long time, by particular properties. These viruses are more resistant to neutralizing antibodies and chemokines. In contrast to cell culture-adapted viruses, patient isolates use different co-receptors for the infection of cells. As patient isolates differ greatly from laboratory-adapted viruses, the collected V2 loop or V3 loop sequence data of the *env* genes of such viruses are to be used for the preparation of the GP120 mixture. Within the framework of the invention, additional base exchanges are possible in regions outside the sequence sections coding for V2 and V3.

According to a particular version of the invention, a protein vaccine is made available which comprises a mixture of GP120 proteins of the human immunodeficiency virus (HIV) which in each case differ from each another in their amino acid sequence in the region of the V3 loop and/or of the V2 loop.

Within the framework of the present invention, during the genetic-engineering preparation of the protein mixture or the protein vaccine, sequence variations are introduced preferably in regions of the V3 loops which lie outside consensus sequences (cf. e.g. M. Schreiber et al., J. Virol. 71 no. 12 (1997) 9198-9205). The consensus sequences are sequence sections which essentially remain both between different virus strains and during the formation of virus variants as a viral illness progress (HIV illness) in the body. Such a section is the sequence Gly-Pro-Gly-Arg-Ala-Phe (GPGRAPH) in the case of the HIV-1 of B sub-type. To the left and right of this sequence unit lie short regions 4-10 amino acids long which can vary greatly. In Fig. 1 this is represented using as example the V3 loop sequence variations of patient isolates.

In addition to variations in the region of the V3 loop, the molecules of the protein vaccine according to the invention can also contain sequence variations in the region of the V2 loop. Changes in the amino acid sequence outside consensus sequence regions are preferred in this case also (see Fig. 2). Furthermore, additional amino acid exchanges are possible in regions outside the V2 and V3 loops.

In connection with the present invention the normal one- or three-character codes are always used to describe the amino acids. When there is variation of the amino acid sequences inside the protein vaccine according to the invention, any desired amino acid exchanges are possible. In each case, however, amino acids of the viral GP120 sequence or of the sequences of the V2 loop and V3 loop are preferably exchanged for amino acids which also occur in other virus

variants at the corresponding sequence positions (see Fig. 2, position of 1997 sequence data in: Human Retroviruses and AIDS, A compilation and analysis of nucleic acid and amino acid sequences, Los Alamos National Laboratory, Los Alamos, NM 87545, U.S.A., Editors: B. Korber, B. Hahn, B. Foley, J.W. Mellors, T. Leitner, G. Myers, F. McCutchan, C. Kuiken).

Within the framework of the present invention, for the first time a protein vaccine is made available which is prepared by genetic engineering and comprises sequence variants of specific viral antigens (proteins). The preparation of the vaccine is described in the following using the V3 loop as an example. It is self-evident to the person skilled in the art that the synthesis principle described below can be transferred to other viruses or viral proteins or parts of their sequences.

In short, a mixture of GP120 proteins with variable V3 sequences which are modelled on the variability of the V3 loop of the GP120 protein of HIV is prepared as a basis for the vaccine.

For this, the sequence coding for the GP120 protein is firstly cloned piecewise into the plasmid pUC 18/19. This takes place through a polylinker exchange so that all interesting sections with monovalent restriction sites are flanked by the possible restriction sites which were previously inserted in the gp120 sequence by silent mutations, and thus can be subsequently cut out and exchanged.

Two homologous nucleic acid oligomers are then prepared with the help of a DNA synthesizer (length approx. 300 base pairs). After hybridization has taken place, the double-stranded DNA fragment forms with the DNA sequence of the V3 loop. Various standard methods from molecular biology can be used for the preparation of the double-stranded V3 loop DNA fragment. In one method, the variables are introduced by inosines and in the case of the corresponding complementary strand the variables are added by a nucleotide mix (AGCT, AGC, AG, ...). In this method, the DNA fragment is prepared by exclusively chemical means. In another method, a combination of chemical and enzymatic DNA synthesis, the variables are introduced by nucleotide mixtures. The oligonucleotides are then hybridized at complementary ends of a length of 20-30 bases, which are not variable. The synthesis to produce the completely double-stranded molecule takes place enzymatically using a DNA polymerase. Both isothermal DNA polymerases (Klenow fragment) or thermostable DNA polymerases (Taq-polymerase) can be used. If Taq polymerase is used, then larger amounts for the cloning of the V3 loop DNA fragment can also be prepared using the polymerase chain reaction. Using these methods, a double-stranded DNA fragment is prepared which is variable at specific positions. This degenerated DNA sequence codes for the corresponding variety of V3 loop amino acid sequences of the GP120 protein mixture to be prepared, the protein vaccine.

The mixture of the synthesized V2 fragments has a PstI cleavage site at the 5' end and a BclI cleavage site at the 3' end. The mixture of the synthesized V3 fragments has a BglII cleavage site at the 5' end and an XbaI cleavage site

at the 3' end. Using the two respective cleavage sites, the fragment mixture is cloned into a vector, e.g. pUC18 delta-env or pUC 18 BstEII-BamHI (Fig. 3). After this cloning a mixture or a pool of plasmid DNAs of the vector pUC 18 forms, which all have the complete BstEII-BamHI env fragment. All plasmid DNAs differ exclusively in the sequence of the env V2 loop or V3 loop. By inserting the V2 or the V3 fragments, the deletion (delta env) of the env gene section in the vector pUC18 is stopped.

This thus-prepared mixture of the plasmid DNA with variable env fragments is transformed into *E. coli* and fermented. *E. coli* amplifies and replicates this mixture of plasmid DNA. All possible variables of these V2/V3 loop plasmids are produced.

This plasmid pool is then isolated. The env fragment is then cut out of the mixture of the plasmid DNA by BstEII and BamHI digestion and cloned direct into the vector for the expression of the viral gp120. This vector has the advantage that the GP120 is also expressed in the eukaryotic cell system.

This pool of BSCenvATG-vector DNA with variable gp120 construct is now transfected into Cos or Chinese hamster ovary cells (CHO cells). These eukaryotes then express this pool of plasmids so that statistically the corresponding protein is translated to each variable and then - depending on which eukaryotes are used - correspondingly glycosylated. The proteins are then harvested, followed by purification to give the finished product (protein mixture or GP120 mixture with variable amino acid sequence).

By way of illustration, the process for producing the protein vaccine according to the invention is represented clearly in Fig. 4.

As already shown, variations in the V2 loop and/or in the V3 loop can be carried out. The variation of the V2 loop can take place according to the same scheme as described for V3.

The preparation of the protein vaccine according to the invention takes place, as described above, via various key constructs, i.e. nucleic acid intermediate steps and DNA constructs, which are essential for carrying out the invention.

As already explained, the nucleic acid sequence coding for the GP120 protein is cloned piecewise into pUC 18/19, the starting point according to the invention being a gp120 sequence in which the repeatedly present restriction sites are firstly modified by silent mutations such that each of the remaining restriction sites still occurs only once in the sequence. This sequence modification is necessary in order to be able to insert the expression cassette, which is necessary for producing the sequence variants, selectively at a quite specific point which is defined by two restriction sites each occurring only once in the sequence. Methods for the introduction of silent mutations in a nucleic acid sequence are well known to a person skilled in the art.

A subject of the present invention is therefore also a nucleic acid sequence which is derived from the env sequence in SEQ ID NO: 1 or a fragment of same, it being modified such that it contains exclusively monovalent restriction sites. Preferably, the modification takes place by the introduction of silent mutations. According to a preferred version of the invention, the nucleic acid sequence has the sequence shown in SEQ ID NO: 9. The sequence of the gene cassette can be modified such that it contains the total env gene or parts of the env gene of virus isolates from patients. Preferably, such a gene cassette is to contain the env sequence of the patient isolate PI-932 (SEQ ID NO: 11). A gene cassette preferred according to the invention based on the PI-932 sequence is shown in SEQ ID NO: 12.

A subject of the present invention is therefore also a single-stranded nucleic acid sequence which contains the region coding for the V3 loop and/or for the V2 loop or fragments or parts of same in which, in the case of the V3 loop, a BglIII-XbaI (247 bp) or also a BglIII-NheI (283 bp) fragment can be exchanged for a modified fragment which carries nucleic acid exchanges or mutations at least 6, preferably at 9 to 20 positions, and in the case of the V2 loop, a PstI-BclI (139 bp) or also a PstI-EcoRI (339 bp) fragment for a modified fragment which carries nucleic acid exchanges or mutations at least 6, preferably at 9 to 20 positions. The nucleotides within a nucleic acid sequence are in each case replaced either by inosine or by a mixture of 2-4 nucleotides. This is explained using the example in Fig. 5. If various amino acids are to be combined at 7 amino acid positions of the V3 loop 21 to give 1152

variants, two nucleotides must be introduced by chemical synthesis at each of 11 nucleic acid positions into the sequence of the single-stranded nucleic acids (oligonucleotides) (Fig. 5).

The single-stranded nucleic acid sequences are - as already mentioned above - hybridized or synthesized to give the double strand.

The present invention therefore furthermore relates to double-stranded DNA which comprises hybrids of the above-mentioned single-stranded nucleic acid sequences, where in each case one nucleic acid sequence (5'-3' oligomer or 3'-5' oligomer) contains inosines at one or more selected positions, and the other nucleic acid sequence (3'-5' oligomer or 5'-3' oligomer) in each case contains two, three or four of the possible nucleotides (adenine, A; thymine, T; guanine, G; cytosine, C) at the corresponding complementary positions during the later hybridization. Sequences are thereby produced in which the four bases are present randomly distributed at the respective positions of the single-stranded nucleic acid sequence (DNA) so that the desired combinations of A, T, G or C are produced for the corresponding desired amino acid codons (a codon is formed from a series of three nucleotides). Because of the variation of nucleotide positions, randomly distributed sequence combinations result. Therefore, e.g. for the combination of (A, C) (ACT) (ACGT) a total of $2 \times 3 \times 4 = 24$ possible DNA sequences form, which code for various amino acids. At the same time, the number of variable positions also determines the number and position of the introduced inosines in the complementary DNA sequence. The calculation

of the heterogeneity of such a construct is shown in Fig. 5.

As, to form the double-stranded DNA, singled-stranded DNA containing inosines is in each case hybridized with oligomers containing A, T, G or C in each case randomly distributed at the corresponding positions, a mixture of double-stranded *gp120*-sequence or of part (fragment) thereof is obtained, the nucleic acid sequences in each case being derived from the *env* sequence (SEQ ID NO: 1 or 12) and the nucleic acid sequences in each case being distinguished from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop. This means that nucleic acid sequences contained in the mixture differ from each other such that they code for a mixture of proteins which in each case contain amino acid sequences which are different from each other in the V2 loop and/or in the V3 loop.

According to a preferred version of the invention, in this way, at least 10^2 , preferably at least 10^3 and according to a particularly preferred version of the invention at least 10^4 sequence variants can be obtained at nucleic acid level (DNA level). This DNA mixture is hereafter called the DNA pool, and in relation to the variants of the *gp120* sequence this is hereafter called the pool with variable *gp120* construct (*gp120* pool).

Within the framework of the present invention, the term sequence variants of a DNA sequence is taken to mean molecules which have a derived nucleic acid sequence vis-à-vis a native viral DNA molecule or a part (fragment) of

same, the variants being distinguished from each other in that at least one nucleotide can be exchanged at any chosen points of the sequence. Preferably, the sequence variants contain several nucleotide exchanges at various points of the sequence, the number and the position of the exchanged nucleic acids essentially depending on the length of the nucleic acid sequence.

Upon the expression of the env-gene variants starting from the produced plasmid-DNA mixture, it is to be expected that, because of the random distribution, all possible DNA sequences are expressed. Therefore, all sequence variants that are possible, on the basis of the pre-set DNA mixture, of the *gp120* molecule or of part of same are then also formed. The heterogeneity of the *gp120* mixture is calculated on the one hand using the heterogeneity of the DNA sequence and the degeneration of the genetic code for the amino acids (cf. also Fig. 5).

The heterogeneity of the plasmid-DNA mixture is demonstrated by DNA sequencing of single clones. To this end, the mixture of the plasmid DNA is transformed in *E. coli* and single clones are randomly selected and their V2 or V3 loop sequence determined. In total, approx. 100-200 different clones are to be sequenced. The statistical distribution of the DNA sequences can be used to calculate the distribution of the whole mixture. The method corresponds very largely to the method of random sampling such as is used as standard for the quality control of the most varied products.

The direct molecular analysis of the heterogeneity of the GP120 protein mixture proves to be somewhat more difficult, as single GP120 molecules cannot be separated from the mixture and identified. This is in the nature of things, as the individual GP120 variants differ from each other in a few amino acids only. Gel electrophoresis separation or mass spectroscopic analysis can establish whether a mixture or a single form of the GP120 molecule is involved. When the GP120 mixture is used e.g. in animals as a vaccine, the immune response induced in the animal is directly dependent on the number and the composition of the GP120 mixture. The immune response of the animal, the neutralizing antibodies can then be examined in virus neutralization tests. To monitor a corresponding variant-overlapping immune response, various patient isolates of HIV-1 can be used in the neutralization test. Preferably, patient isolates are chosen which differ in the ability to use various co-receptors for the infection of the target cells. The neutralization potential of the GP120 mixture then serves as a quality measure.

A subject of the present invention is therefore furthermore a protein mixture which comprises GP120 proteins which contain amino acid sequences which in each case differ from each other in the V2 loop and/or in the V3 loop, the mixture containing, according to one version of the invention, at least 10^2 , preferably at least 10^3 , according to a version of the invention, and, according to a particularly preferred version of the invention, at least 10^4 sequence variants.

As already mentioned earlier, the mixture of double-stranded DNA which can be obtained by hybridization of inosines-containing single-stranded DNA with randomly distributed A-, T-, G- and/or C-containing single-stranded DNA, i.e. the pool with variable GP120 construct, is transformed and fermented in prokaryotic or eukaryotic host cells, preferably *E. Coli*.

A subject of the present invention is therefore plasmids which contain inserted double-stranded DNA, in each case comprising hybrids of the single-stranded, inosine-containing nucleic acid sequence (see above) with the single-stranded nucleic acid sequence containing a mixture of all four nucleotide variants (A, T, G, C) (see above). Furthermore, the invention relates to a vector mixture which contains a mixture of these plasmids, the nucleic acid sequences of the plasmids in each case differing from each other in the region coding for the V3 loop and/or in the region coding for the V2 loop. According to one version of the invention, the vector mixture comprises at least 10^2 , preferably at least 10^3 and according to a particularly preferred version of the invention at least 10^4 of the named plasmids. Depending on which host cells are to be used to express the vectors (plasmids), various expression systems and base vectors well-known to a person skilled in the art can be considered (cf. Methods in Enzymology, Vol. 185, Gene Expression Technology, 1991, Publ. D.V. Goeddel, Academic Press, Inc.). Thus, for the expression in *E. coli*, the plasmid pUC 18/19 is preferred as base plasmid into which the nucleic acid sequence variants are cloned. Within the framework of the present invention, the vector mixtures can thus be expressed either in bacterial host cells such

as *E. coli*, or in eukaryotic host cells, preferably from the group consisting of Cos, CHO or baby hamster kidney cells (BHK cells) or in other host cells.

The present invention thus furthermore relates to *E. coli* host cells or eukaryotic host cells which are transfected with a vector mixture according to the invention.

As host cells are transfected with the DNA pool (the vector mixture), it is to be expected that as great or at least approximately as great a number of differently transformed host cells forms as there are sequence variants present in the DNA pool. Upon the preparation of the vaccine, special attention is to be paid to the yields during the individual cloning steps. The yields upon preparation of the double-stranded DNA fragments, the ligation of the V3 loop and V2 loop DNA fragments into the *env* gene and the transformation or the preparation of bacteria and cell clones must be such that the heterogeneity that is to be achieved of the mixture is not restricted. This is to be illustrated using an example. If, e.g. a number of balls are to be removed from a quantity of balls which differ on the basis of two colours, but there is to be a 99.9% probability that both colours are present in this selection, approx. 13 balls would have to be removed (G. Schreiber, Ein kombinatorisches Problem aus der Genetik; Bio-engineering 1988, 2: 32-35). Transferred to the cloning steps of the HIV vaccine, this means: If a heterogeneity of approx. 6000 virus variants is to be produced, 13 times as many clones would have to be produced in each individual cloning step. Thus, a gene bank of approx. 80,000 clones must be prepared for the V3 loop. Starting from this gene bank, the

expression vectors for the transfection of the CHO cells are then prepared. During transfection of the CHO cells, consequently, approx. 80,000 transfections must then be achieved. Such a mixture of transfected CHO cells then produces the desired quantity of 6000 different gp120 variants.

The present invention also relates to a process for the preparation of a nucleic acid sequence coding for a viral protein (expression cassette) in which the sequence is so modified, or preferably so many silent mutations are introduced, that it then contains at least two and preferably exclusively monovalent restriction sites. Particularly preferably, the sequence still contains only monovalent restriction sites. Preferably, the protein coded by the nucleic acid sequence is GP120, the env wild-type sequence coding for the GP120 being varied by silent mutations.

A further subject of the present invention is a process for the preparation of the vector mixture according to the invention, which preferably contains a mixture of plasmids, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop, the plasmids according to the invention being ligated into a (base) vector which can be expressed in host cells (preferably *E. coli*, Cos, CHO or BHK cells). The (base) vector is preferably the pUC 18, pUC 19 or BSCenvATG vector.

Within the framework of the present invention, a process is furthermore made available for the preparation/production

of host cells, preferably chosen from the group consisting of *E. coli*, Cos, BHK or CHO cells, in which the host cells are transformed with a vector mixture according to the invention, which contains a mixture of plasmids the nucleic acid sequences of which in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop.

Finally, according to the invention, a process is for the first time made available for the preparation of a DNA vaccine, in which the process according to the invention for the preparation of the vector mixture is carried out, the plasmids according to the invention expressing the mixture of the *gp120* proteins after application in host cells (e.g. human and animal monocytes). The DNA vaccine can optionally be formulated with pharmaceutically compatible auxiliaries and/or carriers for application and after administration in the organism, make possible the production of the sequence variants of viral proteins.

The present invention thus also relates to a pharmaceutical composition or a DNA vaccine which codes for a mixture of structurally different virus proteins, the vaccine containing a mixture of sequence variants of a viral DNA molecule or of part of same, i.e. of DNA molecules the nucleic acid sequences of which differ from each other in the region coding for the protein, a part or a fragment of same. By the term "structurally different virus proteins" is meant according to the invention those proteins the amino acid sequences of which are derived from the wild-type sequence of the corresponding virus protein, the amino acid sequences deviating from each other insofar as they

different from each other and compared with the wild-type sequence through one or more exchanged amino acids at identical or different positions of the sequence. As already mentioned, the nucleic acid sequences in the vaccine code preferable according to the invention for a mixture of structurally different GP120 proteins of HIV (sequence variants of GP120), a vaccine being particularly preferred which contains a mixture of DNA molecules, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop of *gp120* of the HIV-1. In this context, reference is also made to Fig. 3 in which known structural differences or variations in the V3 loop of GP120 are shown.

The DNA vaccine according to the invention is of particular importance within the framework of a gene therapy.

Finally, according to the invention, a process for the preparation of a pharmaceutical composition or a protein vaccine is for the first time made available in which the host cells according to the invention, i.e. host cells which are transformed with a vector mixture according to the invention, the vectors in each case containing plasmids, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop, are cultivated under conditions which allow the expression of the mixture of viral protein sequence variants. The host cells are preferably bacterial host cells, such as *E. coli*, or eukaryotic host cells, preferably from the group consisting

of Cos, Chinese hamster ovary (CHO) or baby hamster kidney cells (BHK cells).

Through the present invention, it is thus possible to make available a vaccine comprising variable GP120 proteins which, in the prophylactic and/or therapeutic treatment of a HIV infection or AIDS

1. activates the immune system,
2. induces the formation of HIV-1-neutralizing antibodies,
3. prevents the loss of neutralizing antibodies and
4. in which monitoring of the stimulation of the GP120-specific immune response is taken over by the active ingredient.

This is preferably carried out in that, even before the loss of HIV-neutralizing antibodies associated with the outbreak of AIDS, sufficient antigens are available which, because of their variety of different amino acid sequences (i.e. their sequence variations) are in a position to induce the formation of those HIV antibodies which normally, i.e. without corresponding prophylaxis according to the present invention, are lost or experience a reduction in concentration as the illness progress. In addition to a preventative treatment, the present invention also relates to therapeutic treatment, the GP120 mixture (protein vaccine) being used, to counter a reduction or a loss of HIV-neutralizing antibodies by activation of the immune system and induction of the formation of new or additional HIV-1-neutralizing antibodies.

According to the invention, the mixture of structurally different viral proteins which are sequence variants of a viral protein (preferably of GP120) or of part of same is therefore used for the preparation of a vaccine for the prevention and or therapy of a viral infection in humans. Furthermore, the present invention relates to the use of a mixture of DNA molecules which code for sequence variants of a viral protein or of part of same for the preparation of a vaccine for the prevention and/or therapy of a viral infection in humans. According to a preferred version, the invention relates to the use for the preparation of a vaccine for the prevention and/or therapy of a HIV infection in humans.

Within the framework of the present invention, the virus infection can be any infection in which self-replicating virus variants form in the course of the illness, where those amino acid sequence sections or nucleic acid sections coding for the latter can be considered according to the invention as selected protein sequences or protein-coding nucleic acid sequences, in which, in the course of viral illnesses, variants, i.e. sequence variations, are always or often observed. Preferably, in the present case, these are sequences variants of the GP120 molecule (at protein level) or of the region coding for GP120 (at DNA level) or parts of same, in particular sequence variants in the V3 loop or in the V2 loop, preferably both in the V3 and in the V2 loop.

According to the invention, pharmaceutical compositions or vaccines are consequently made available which, in the immunoreconstitutive treatment of those suffering viral

infection, activate the immune system in such a manner that the naturally acquired immune defence is regenerated against the virus and restimulated to thus prevent the virus variants replicating in the patient from multiplying. According to a preferred version of the invention, vaccines for the immunoreconstitutive treatment of those infected by HIV-1 are made available for the first time.

Within the framework of the present invention, the vaccines can either be formulated or administered per se, i.e. as DNA and/or protein mixtures without further additives, or together with further active ingredients, such as e.g. immunostimulants such as interleukin-2, CC and CXC-Chemokine, and/or pharmaceutically compatible auxiliaries and carriers. As a rule, the vaccines according to the invention are formulated for intravenous application, such as e.g. by the intravenous, intramuscular, subcutaneous routes. In addition, the vaccines can be formulated for oral, mucosal (intravaginal, intrarectal) and transdermal application.

The present invention has the following advantages:

Through a selective genetic manipulation of the nucleic acid sequence, with the help of a gene cassette, any desired variants of a pathogen or of a gene of the exciter can be prepared. The variations preferably affect the regions against which the neutralizing antibodies or cytotoxic T cells are formed. An inoculant in the form of a mixture of variants of the pathogen or of variants of the corresponding antigens of a pathogen can thereby be prepared. The advantages of this concept in relation to HIV

as a pathogen lie in the preparation of a mixture of HIV-GP120, the external membrane protein of HIV, against which the virus-neutralizing immune response of humans is directed. As, in the precise case of the HIV illness, each patient is exposed to a variety of HIV variants which all differ in the sequence of the GP120 protein, the use of a variety of GP120 variants as inoculant is of particular advantage, this GP120 mixture being able to be used both for the immunization of normally healthy persons and for the therapy of persons already infected by HIV. In the case of immunization with the GP120 mixture, as broad as possible an immune response against as many virus variants as possible in humans is induced, which in the ideal case protects against all HIV variants. In the case of therapy with the GP120 mixture, the loss of neutralizing antibodies and the loss of the HIV-specific cellular immune response can be combated.

According to the invention, it is furthermore possible for the first time to prepare *gp120* clones which, in the nature or variety of the observed sequence variations, correspond to plasma isolates of those infected with HIV or ill with AIDS (cf. M. Schreiber et al., J. Virol. 68 No. 6 (1994) 3908-3916). In this context, in contrast to various approaches followed to date, no isolated GP120 molecule or an antigenic fragment of same is used for immunization, but a protein mixture which consists of a variety of sequence variants which are characterized by a complete GP120 sequence which in addition contain the correct tertiary structure corresponding to the natural plication of the GP120 molecule. The conformation of the virus variants made available in the vaccine is important insofar as sequence

and structure variants are thereby provided which, with the GP120 variants detectable in the course of the viral illness, as great as possible a similarity with regard to the observed sequence variations and also the conformation required for the bonding to CD4 and the co-receptor is achieved, whereby an effective immune stimulation can be achieved.

Through the invention, a mixture of GP120 proteins is provided which represents the protein vaccine. At the same time, a mixture of the genes coding for this protein mixture is provided through the invention. These genes can be transferred into a vector which is suitable for direct application in humans (DNA vaccine) (Ulmer et al., 1995 Ann NY Acad Sci 772:117-125; Donnelly et al., 1995 Ann NY Acad Sci 1995 772:40-46). A DNA vaccine has the advantage that the heterogeneity of the mixture of the DNA vectors can be greater than that of the mixture of the recombinant GP120 proteins. It is easier to produce a high heterogeneity of a DNA vector mixture. Such a DNA vaccine would cover a much broader spectrum of HIV variants. From the technical point of view, the preparation of the DNA mixture is easier. A precondition for the DNA vaccine is a *gp120* expression vector which carries the two cleavage sites BstEII and BamHI. This makes possible the conversion of the V3 loop and V2 loop variables env gene fragments into such a vector.

With regard to the GP120 proteins, it is to be borne in mind that 5 potential points for N-glycosylation are located in the region of the V3 loop. The sugar groups protect the V3 loop against recognition by neutralizing

antibodies. Virus variants which contain a glycosylated V3 loop are more poorly neutralizable than virus variants in which glycosylation is incomplete. As virus variants with a complete V3 loop glycosylation can escape the neutralizing immune response, they are important for the transfer and for the establishment of the infection. In the course of the illness, if a part of the neutralizing antibody response is lost, virus variants whose V3 loop is not completely glycosylated establish themselves. As the V3 loop is no longer covered by sugar groups, these virus variants replicate faster than the completely glycosylated V3 loop mutants. According to the invention, it is therefore advantageous to take the varying glycosylation of the GP120 V3 loop into account when constructing the GP120 vaccine.

Within the framework of the present invention, the principle represented with reference to HIV can be transferred to other viruses which also form variants in the course of the viral illness. In particular, vaccines can be provided against a variety of viruses which induce a broad immune response against many virus variants in humans which in the ideal case protects against all variants. In the case of therapy with such vaccines, the loss of neutralizing antibodies and the loss of the HIV-specific cellular immune response can be effectively combated. In this context, it is furthermore possible or even advisable, with pathogen types (viruses) which form several sequence variants of several proteins in the course of an illness, to provide several gene cassettes according to the invention each of which contains the nucleic acid sequence coding for variants of each of these proteins or fragments

of same, in order to combat the loss of neutralizing antibodies against all conceivable virus variants.

Within the framework of the present invention, the advantages of a protein vaccine and a DNA vaccine can be advantageously combined in order to increase the success of a preventative and/or therapeutic treatment of a viral illness. A pharmaceutical composition is thus furthermore provided for the prevention and/or therapy of a viral infection, which comprises a protein mixture and a nucleic acid mixture, the protein mixture comprising sequence variants of a viral protein or of part of same, and the nucleic acid mixture comprises DNA molecules which code for sequence variants of a viral protein or of part of same. In particular, the pharmaceutical composition is a combination preparation which comprises both a protein mixture comprising one of the above-named sequence variants of the GP120 protein and a nucleic acid mixture derived from an above-named env sequence in SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same.

The present invention is explained in the following, using examples, figures and a sequence protocol.

EXAMPLES

1. General description of the preparation process
 - 1.1 Cloning of V3 loop-coding oligonucleotides
 - 1.2 Cloning of the V3 env variants
 - 1.3 Analysis of variability
2. Material and methods
 - 2.1 Abbreviations
 - 2.1.1 General abbreviations
 - 2.1.2 Nucleic acids
 - 2.2 Bacterial strains
 - 2.3 Plasmids
 - 2.4 Enzymes
 - 2.5 Chemicals
 - 2.6 Oligonucleotides
 - 2.7 Molecular weight standards
 - 2.8 Reagent kits used
 - 2.9 Media
 - 2.10 Sterilization of solutions
 - 2.11 Cultivation and storage of bacteria
 - 2.12 Preparation of competent cells
 - 2.13 Transformation into E. coli
 - 2.14 Plasmid-DNA preparation
 - 2.15 Separation of DNA in agarose gels
 - 2.16 Purification of DNA from agarose gels
 - 2.17 Separation of DNA in polyacrylamide gels
 - 2.18 Cleaving of DNA
 - 2.19 Ligation of DNA
 - 2.20 DNA sequencing

FOOTED" 6516660

- 2.21 Preparation of double-stranded DNA with the help of oligonucleotides
- 2.22 Transfection of COS cells and CHO cells
- 2.23 Chromatographic purification of the GP120 mixture
- 2.24 Preparation of the DNA vaccine

1. General description of the preparation process

1.1 Cloning of a mixture of V3 loop *gp120* variants

The oligonucleotides are synthesized as described under 2.6. As V3 loop sequence, a mutated sequence of the HIV-1 patient isolate F1-01 (M. Schreiber et al., J. Virol. 68 no. 6 (1994) 3908-3916) is given by way of example. Any other sequence or mixture of sequences is also possible. To clone many different variants of a V3 loop sequence, mixtures are used instead of pure nucleotide components at specific positions of the sequence. In this way, a mixture is obtained of oligonucleotides which all differ in the sequence and thus code for different V3 loops. Such oligonucleotide mixtures are also called oligonucleotides with degenerated sequences. Starting from chemically synthesized oligonucleotides with degenerated sequences, the region coding for the various V3 loops is shown.

An oligonucleotide, in reading-frame orientation (forward), is synthesized for the *env*-sequence section from the BglIII cleavage site to the first degenerated position. A second oligonucleotide, in complementary orientation (reverse), is synthesized according to the sequence from a position which lies 15 bases before the beginning of the variable region (method 2.6). Through overlapping of the 3' region along a

length of 15 bases in total, the hybridization of both oligonucleotides is carried out. In a subsequent reaction with DNA polymerase (e.g. Taq DNA polymerase or Klenow fragment), a completely double-stranded DNA molecule forms from the two hybridized oligonucleotides (method 2.21).

The thus-obtained DNA mixture is digested with restriction endonucleases BglII and XbaI. The cloning of the DNA mixture is carried out in the expression vector Δ V3 pBSCenvV3 cleaved with BglII and XbaI (methods 2.15 and 2.16).

This vector contains the coding sequence of the *gp160* of the HIV-1 strain NL4-3 (IIIB). The NL4-3 *env* gene was manipulated such that it possesses the restriction cleavage sites BglII and XbaI as well as ApaLI, PstI and BclI once each only. The region coding for the V3 loop, which lies between the cleavage sites BglII and XbaI, was removed and replaced by a 15 base pair sequence, whereby an analytical cleavage site was introduced for the enzyme AscI. The BglII- and SbaI- cleaved V3 loop DNA mixture is cloned into this vector (method 2.19). The loss of the AscI cleavage site in the finished V3 loop pBSCenvV3 vector can be used for the selection of the V3 loop coding clones (method 2.18).

The thus-formed plasmid mixture is transformed in DH5 α bacteria (method 2.12), a transformation rate of $>10^5$ having to be achieved. A gene library of V3 loop fragments with a size of approx. 10^5 clones is thereby obtained. The mixture is to be analyzed by DNA sequencing (see 1.3). With the help of this process, a mixture of clones is obtained,

which all code for different V3 loop variants of the GP120 protein. This mixture of vectors serves as starting product for the preparation of the protein mixture for use as inoculant and as starting product for use as DNA vaccine.

For the preparation of the GP120 protein mixture, the pBSCenvV3 expression vectors are transfected into COS cells and CHO cells (method 2.22). The purification of the different GP120 proteins, the active ingredient mixture, is carried out according to method 2.23, as described in the literature.

1.2 Cloning of a mixture of V2 loop *gp120* variants

Variants of the V1 loop and V2 loop are prepared in the same way. The expression vector $\Delta V3$ -pBSCenvV3 has three additional restriction cleavage sites for this. The variation of the V1 loop takes place through cloning into the *Apa*LI and *Pst*I cleavage sites. The variation of the V2 loop takes place through cloning into the *Pst*I and *Bcl*I cleavage sites.

1.3 Analysis of variability

The analysis of the V3 loop mixtures takes place through DNA sequencing (method 2.20). The statistical selection of approx. 100-200 clones whose V1, V2 and V3 loop sequences are to be determined, is planned. If all these clones are different, the heterogeneity of the gene library and thus also the heterogeneity of the protein mixture can be measured with the help of a statistic calculation.

2. Material and methods

2.1 Abbreviations

2.1.1 General abbreviations

µg microgram
µl microlitre
µmol micromole
APS ammonium persulphate
bp base pairs
dNTP desoxynucleoside triphosphate
DNA desoxyribonucleic acid
DTT dithiothreitol
EDTA ethylenediaminetetraacetic acid
g gram
GP glycoprotein
h hour
HIV human immunodeficiency virus
IPTG isopropyl-β-D-thiogalactopyranoside
MOPS 3-morpholinepropanesulfonic acid
OD optical density
PAGE polyacrylamide gel electrophoresis
PCR polymerase chain reaction
RT room temperature
TEMED N,N,N',N'-tetramethylethylenediamine

2.1.2 Nucleic acids

A adenine
C cytosine
G guanine
T thymidine

2.2 Bacterial strains

Escherichia coli DH5 α : F-, endA1, hsdR17,
(rk-mk+), su-pE44, recA1,
 λ -, gyrA96, relA1, Φ 80d lac
z Δ M15

2.3 Plasmids

pBSCenvATG own design, the sequence is given in
SEQ ID NO: 10.

2.4 Enzymes

Restriction enzymes MBI-Fermentas, Gibco-BRL, Biolabs
DNA polymerases MBI-Fermentas, Gibco-BRL
T4-DNA ligase MBI-Fermentas

2.5 Chemicals

[α -35S]dATP	Amersham Life Science
Agarose, ultra pure	GIBCO BRL
Ammonium persulphate	Merck
Ampicillin	US Biochemical
Bacto-Agar	Becton Dickinson
Bacto-Trypton	Becton Dickinson
Boric acid	Merck
Bromphenol blue	Merck
Calcium chloride	Merck

Desoxyribonucleotides	MBI-Fermentas
Dithiothreitol	Biotechnik, St. Leon-Rot
Glacial acetic acid	Merck
Ethidium bromide	Sigma
Ethylenediaminetetraacetic acid	Merck
Glycerine	Merck
Urea	ICN Biomedicals
Yeast extract	Becton Dickinson
Potassium chloride	Merck
Potassium dihydrogen phosphate	Merck
Magnesium chloride	Merck
Acrylamide mix	Roth
Sodium acetate	Merck
Sodium chloride	Merck
Disodium hydrogen phosphate	Merck
Sodium dihydrogen phosphate	Merck
2-propanol	Merck
Sigmacote (chlorinated polysiloxane)	Sigma
N,N,N',N'-tetramethylethylenediamine	Merck
Tris(hydroxymethyl)-aminomethane	GIBCO BRL

2.6 Oligonucleotides

All oligonucleotides listed were prepared with the ExpediteTM nucleic acid synthesis system of PE Biosystems (Weiterstadt). Oligonucleotides are used for the cloning of

env genes which differ from each other only in the sequence for the V3 loop. The sequences of the oligonucleotides are listed taking as an example the cloning of the V3 loop region for the HIV-1 patient isolate F1-01.

V3 loop: for cloning into pNL4-3/BglIII-NheI, F1-01,
forward:

5'-AAG ATG TAG TAA TTA GAT CTG CCA ATT TCA CAG ACA ATG CTA
AAA CCA TAA TAG TAC AGC TGA ACA CAT CGT TAG AAA TTA ATT GTA
CAA GAC CCA ACA ACA AT ACA-3' (SEQ ID NO: 3)

V3 loop: for cloning in pNL4-3/BglIII-NheI, F1-01:

5'-TTT TGC TCT AGA AAT GTT ACA ATG TGC TTG TCT TAT GTC TCC
TGT TGC AGC TTC TGT TGC ATG AAA TGC TCT CCC TGG TCC GAT ATG
GAT ACT ATG-(GA) (AT) (GATC) TTT TCT TGT ATT GTT GTT GGG-3'
(SEQ ID NO: 4)

For sequencing of V3 loop clones, the oligonucleotides 7010, 7011 and the M13 standard primer were used (M13, M13r).

Primer 7010: 5'-CCA TGT ACA AAT GTC AG-3' (SEQ ID NO: 5)

Primer 7011: 5'-AAA ACT GTG CGT TAC AA-3' (SEQ ID NO: 6)

M13-Primer: 5'-GTA AAA CGA CGG CCA GT-3' (SEQ ID NO: 7)

M13r-Primer: 5'-CAG GAA ACA GCT ATG AC-3' (SEQ ID NO: 8)

2.7 Molecular weight standards

1 kb conductor MBI-Fermentas

100 bp conductor MBI Fermentas

2.8 Reagent kits used

T7 Sequencing-Kit	Pharmacia
Qiaquick PCR purification Kit	Qiagen
Qiagen Plasmid Kit	Qiagen

2.9 Media

YT Medium:

10g trypton
5g yeast extract
5g NaCl

dYT medium:

16g trypton
10g yeast extract
5g NaCl

dYT agar plates:

10g trypton
5g yeast extract
5g NaCl
15g agar

The quoted quantities relate in each case to 1000 ml deionized water. The mixtures were autoclaved at 121°C and 1.5 bar for 20 mins. Should the media contain antibiotics or other heat-sensitive reagents, the corresponding quantities were sterile-filtered and added after cooling of the medium.

Ampicillin 3.3 ml/l (60 mg/ml)

IPTG	3 ml/l (100 mM)
xgal	3 ml/l (2% in DMF, do not filter)

2.10 Sterilization of solutions and apparatus

Solutions and media as well as pipette tips, Eppendorf vessels and glass apparatus were autoclaved for 20-40 mins at 121°C and 1.5 bar. Heat-sensitive solutions such as e.g. antibiotics and IPTG solutions were sterile-filtered.

2.11 Cultivation and storage of bacteria

Bacteria cultures were in each case inoculated with a single colony. To isolate a single colony, cells of a liquid culture were spread on an agar plate. After incubation at 37°C overnight, isolated colonies have grown. For the short-term preservation of the bacteria, the agar plates were sealed with parafilm and stored at 4°C. For a long-term storage of bacteria strains, 0.75 ml of a YT-overnight culture were mixed with 0.25 ml sterile glycerine, shock-frozen in liquid nitrogen and stored at -70°C.

2.12 Preparation of competent cells

100 ml YT-medium were inoculated with 100 µl of an overnight culture. The bacteria were incubated until an $OD_{550}=0.4$ at 37°C was reached in the shaking incubator and then centrifuged for 8 mins at 4°C and 1000g. All subsequent work was carried out on ice using pre-cooled vessels and 4°C cold solutions. The cells were resuspended in 50 ml 50 mM $CaCl_2$ and incubated for 30 mins on ice. After

renewed centrifuging, the bacteria were taken up in 2.5 ml sterile TFBII buffer and divided into portions of 100 μ l each. The 100- μ l portions of the competent cells were able to be, insofar as they were not required for an immediate transformation, stored in liquid nitrogen at -70°C after shock-freezing.

TFBII buffer:

10 mM	MOPS pH 7.0
75 mM	CaCl_2
10 mM	KCl
15%	glycerine

2.13 Transformation of *E. coli*

(Hanahan, J. Mol. Biol. 166 (1983) 557-580)

A 100 μ l portion of competent cells was thawed in an ice bath and incubated with 1-100 ng plasmid-DNA for 30 mins at 0°C . The transformation mixture was then incubated for 1 min at 42°C . 1 ml YT-medium was then added and agitated for an hour at 37°C . To make possible a selection of the transformed cells, 100-500 μ l of the mixture were spread onto antibiotic-containing YT agar plates. After incubation overnight at 37°C , only colonies which carry the plasmid-coded resistance gene have grown. Upon a transformation of cells which do not contain any functional β -galactosidase (lacZAM15 mutation, e.g. DH5 α), vectors such as the pUC used make possible a direct selection (*blue white screening*) of recombinant bacteria via a plasmid-coded β -galactosidase. For the blue-white selection, the bacteria were spread onto Amp/IPTG/Xgal-YT agar plates. The substrate Xgal is cleaved into a blue dye by the β -

galactosidase. Because of the destruction of the reading frame of the lacZ gene by the insertion of a foreign DNA fragment into the lacZ gene, white colonies are produced. On the other hand, blue colonies mean that the lacZ gene has remained functional during cloning and ligation and no DNA was inserted.

2.14 Plasmid DNA preparation

(Qiagen, Hilden)

To isolate plasmid-DNA from *E. coli*, the QIAprep Spin Miniprep Kit from Qiagen was used. The DNA preparation took place according to the manufacturer's instructions (QIAprep Plasmid Handbook 03/95). Plasmid-containing *E. coli* bacteria were inoculated in dYT-amp medium and agitated overnight at 37°C. Three ml of the *E. coli* overnight culture were used for the plasmid isolation. The bacteria were harvested (1 min, 14,000 rpm, Heraeus centrifuge) and taken up in 250 µl P1 buffer. The bacteria were decomposed through alkaline lysis (250 µl, 0.2N NaOH/1%-SDS). The mixture was neutralized through the addition of 3M potassium acetate, (350 µl, pH 5.5). The purification of the plasmid DNA is based on the selective bonding of plasmid DNA to DEAE anion exchanger columns. With the chosen salt concentrations and pH conditions, the plasmid DNA bonds to the DEAE matrix. The DEAE matrix was washed (750 µl, PE buffer, Qiagen). The plasmid DNA was then eluted with 50 µl H₂O and stored at -20°C.

To obtain plasmid DNA in a larger quantity (approx. 100 µg), 25 ml overnight culture were used. To purify the DNA, the QIAGEN Plasmid Midi Kit was used. The purification is

based in this case on the principle described under the "QIAprep Spin Miniprep Kit" and was carried out according to the manufacturer's instructions (QIAGEN Plasmid Purification Handbook 01/97).

2.15 Separation of DNA in agarose gels

The separation of DNA took place through gel electrophoresis in agarose gels. Depending on the size of the fragments examined, 0.8-2% agarose were dissolved by boiling-up in TBE buffer (for analytical agarose gels) or TAE buffer (for preparative agarose gels). After cooling to approx. 60°C, the gel liquid was mixed with 1 µl ethidium bromide (10 mg/ml) and poured into a prepared gel bed with inserted comb. The completely cooled gel was coated in an electrophoresis chamber with TBE or TAE buffer and the comb removed. The DNA samples were mixed with 1/5 vol. coating buffer and pipetted into the sample bags. The fragments were separated at 5-10 volt/cm gel length for 0.5-2 h. For detection, the gel was photographed after the electrophoresis in UV transmitted light. The molecular weights of the DNA bands were measured compared with following DNA molecular weight markers.

TBE buffer:

100 mM	Tris
100 mM	boric acid
3 mM	EDTA

TAE buffer:

40 mM	Tris
2mM	EDTA

0.114% glacial acetic acid

Coating buffer:

0.25% bromphenol blue
0.25% xylene cyanol
30% glycerine
50 mM EDTA

2.16 Purification of DNA from agarose gels

For the extraction of the DNA from agarose gels, the "QIAquick Gel Extraction Kit" (Qiagen, Hilden) was used. The buffer conditions were selected such that the nucleic acids bond to the silica membrane of the columns, whilst low-molecular-weight bacteria components pass through the membrane (method 2.15). The process was carried out according to the "Qiaquick Gel Extraction Kit Protocol" of the QIAquick Spin Handbook 07/97. The purified DNA was in each case eluted from the silica membrane with 50 μ l H_2O .

2.17 Separation of DNA in polyacrylamide gels

The separation of radioactively marked DNA fragments for the DNA sequencing was carried out using denaturizing polyacrylamide gels. Two purified glass plates were cleaned of fatty residues with ethanol and coated with 1 ml Sigmacote per plate to obtain a hydrophobic surface. Through the coating, a later tearing of the gel upon removal from the glass plate was to be avoided. Spacers were placed between the two glass plates, and served at the same time for lateral sealing. The whole apparatus was fixed with several clamps. To prepare the sequence gel, 21

g urea were dissolved in approx. 20 ml water. After the addition of 7.5 ml acrylamide mix and 5 ml 10x TBE buffer (see 2.15), the mixture was made up to 50 ml with water. The polymerization of the gel was started by the addition of 300 μ l APS and 100 μ l TEMED. Immediately afterwards, the gel solution was poured into the prepared gel apparatus and the pocket bottom shaped by inserting the flat back of the saw-tooth comb. The gel was stored horizontally at room temperature until polymerization was complete. After the insertion of the gel into the gel chamber, the latter was filled with TBE buffer. By rotating the saw-tooth comb, the deposit pockets were formed. A 20-min pre-run was carried out in order to heat the gel to the optimal operating temperature. The pockets were washed thoroughly with TBE buffer and then filled with 4-5 μ l of the samples. The electrophoresis took place at 2 kV (150 watt) for approx. 2-3 h. The gel was removed from the gel chamber after the electrophoresis was ended and one of the glass plates raised carefully. By laying-on and light pressing, the gel was transferred onto Schleicher & Schuell paper (Whatman, England). After drying at 80°C under vacuum, the gel was exposed on an X-ray film (Kodak BioMax MR-1, Sigma Deisenhofen) for 1-3 days at RT.

Acrylamide mix:

- 40% polyacrylamide
- 0.8% bisacrylamide

2.18 Cleaving of DNA

Restriction endonucleases recognize and hydrolyze enzyme-specific palindromic DNA sequences of mostly 4-8

nucleotides' length. Upon hydrolysis, depending on the type of enzyme, blunt ends or overhanging ends of single-stranded DNA form. For a restriction digestion, 0.1-60 µg DNA were incubated for 2-5 h at 37°C with 2-120 units of a restriction enzyme in the buffer indicated by the manufacturer. The total volume was between 10 µl for analytical and 300 µl for preparative mixtures. For restrictions with two different enzymes, a buffer was chosen in which both enzymes have a sufficient activity, e.g. the EcoRI buffer for an EcoRI/BamHI digestion, or the conditions for the second enzyme were set after incubation with the first enzyme.

2.19 Ligation of DNA

DNA ligases catalyze the linking of DNA molecules, consuming NAD⁺ or ATP through formation of a phosphodiester bond between a free 5' phosphate group and a 3' hydroxyl group. A three- to five-times excess of DNA fragment was incubated overnight at 12°C in ligation buffer with 100-500 ng cleaved plasmid and 5 units T4-DNA ligase as well as 10 nmol ATP. Only ligations of cohesive ends were carried out.

Ligation buffer:

40 mM	Tris-HCl pH 7.8
10 mM	MgCl ₂
10 mM	DTT
0.5 mM	ATP

2.20 DNA sequencing

(Sanger et al., Proc. Natl. Acad. Sci. USA 74 (1977) 5463-5467) The sequencing of DNA was carried out according to the chain-break method. Double-stranded plasmid DNA serves as substrate. Double-stranded DNA can be synthesized from single-stranded DNA starting from an oligonucleotide primer in the presence of dNTPs through a DNA polymerase. If a small proportion of dideoxynucleotides (ddNTPs) is contained in the nucleotide mixture, this leads to a randomly distributed incorporation of the ddNTPs, as the DNA polymerase cannot distinguish between dNTPs and ddNTPs. The incorporation leads to a breaking of the double-stranded synthesis, because of the absence of the 3'-hydroxyl group of the ddNTPs and, as it takes place statistically distributed, to DNA single strands of different lengths. As the synthesis mixture is divided in four and only one of the four ddNTPs is present in each sample, base-specific chain breaks result in the respective mixtures. To mark the synthesized single strand, α -³⁵S-dATP is added to the reaction. After denaturing and separation of the DNA via polyacrylamide gel electrophoresis, the bands can be autoradiographically detected and the sequence of the DNA strand read off direct.

The sequencings were carried out with the T7-Sequencing-Kit (Pharmacia) according to the manufacturer's instructions. Either the M13-Universal-Primer (Pharmacia) or Primer M13r was used as oligonucleotide primer. 32 μ l (approx. 2 μ g) of purified double-stranded plasmid DNA were denatured with 8 μ l 2 M NaOH for 10 mins at RT. After the addition of 7 μ l 3 M sodium acetate, pH 4.8, 4 μ l H₂O and 120 μ l -20° ethanol, the DNA was precipitated for 20 mins at -70°C. The DNA was isolated by centrifuging (15 mins, 4°C), washed twice with

-20° cold 70% ethanol and dried in a vacuum centrifuge. The DNA sediment was then resuspended in 10 µl H₂O and after the addition of 2 µl annealing buffer and 2 µl primer solution (10 pmol in water) hybridized with the oligonucleotide primer at 65°C for 5 mins, 37°C for 10 mins and 5 mins at RT. Directly afterwards, for the purpose of primer elongation and radioactive marking, 3 µl labelling mix, 1 µl α-³⁵S-dATP (1000 Ci/mmol, 10 µCi/µl) and 2 µl T7 polymerase solution (diluted 1:5 with enzyme dilution buffer) were added and incubated for 5 mins at RT. In each case, 4.5 µl of this mixture were placed onto a *MicroSample Plate* (Greiner) preheated to 37°C in which 2.5 µl each of the four different dNTP/ddNTP mixes were present. After five minutes' incubation at 37°C, which served for further elongation and base-specific termination, the reactions were ended by the addition of 5 µl stop solution. Before application to the polyacrylamide gel, the samples were denatured for 2 mins at 80°C. The samples were stored at -20°C.

Annealing buffer:

1 M	Tris-HCl pH 7.6
100 mM	MgCl ₂
160 mM	DTT

Labelling mix:

1.375 µM	dCTP
1.375 µM	dGTP
1.375 µM	dTTP
333.5 mM	NaCl

Enzyme dilution buffer:

20 mM	Tris-HCl pH 7.5
5 mM	DTT
100 µg/ml	BSA
5%	glycerine

Stop solution:

10 mM	EDTA
97.5%	formamide
0.3%	bromphenol blue
0.3%	xylene cyanol

A-mix:	840 µM dCTP	C-Mix:	840 µM dATP
	840 µM dGTP		840 µM dGTP
	840 µM dTTP		840 µM dTTP
	93.5 µM dATP		93.5 µM dCTP
	14 µM ddATP		14 µM ddCTP
	40 mM Tris-HCl pH 7.6		40 mM Tris-HCl pH 7.6
	50 mM NaCl		50 mM NaCl
G-Mix:	840 µM dATP	T-Mix:	840 µM dATP
	840 µM dCTP		840 µM dCTP
	840 µM dTTP		840 µM dGTP
	93.5 µM dGTP		93.5 µM dTTP
	14 µM ddGTP		14 µM ddTTP
	40 mM Tris-HCl pH 7.6		40 mM Tris-HCl pH 7.6
	50 mM NaCl		50 mM NaCl

2.21 Preparation of double-stranded DNA using oligonucleotides

The oligonucleotides were hybridized at 60°C for the preparation of the DNA mixture. 100 pmol were used per

nucleotide. 1 pmol of the hybridized sample was used for the PCR or 100 pmol for the Klenow reaction.

The following cycles were carried out for the PCR:

At the start:	10 min	94°C
30 cycles:	1 min	94°C
	1 min	45°C
	1 min	72°C
At the end:	10 min	72°C

1 pmol hybridized oligonucleotide was used as DNA *template* for the PCR. PCR primers were used for all PCR reactions carried out in a final concentration of 0.1 pmol/ μ l. 1 unit of the Taq polymerase was used per 50 μ l of the reaction mixture. The concentration of the dNTPs in the PCR mixture was set at 0.1 mM.

To prepare the DNA mixture with the help of the Klenow reaction, 100 pmol of the hybridized oligonucleotides was dissolved in Klenow buffer (50 mM Tris-HCl, pH 8.0; 50 mM $MgCl_2$, 10 mM DTT, 0.05 mM dNTP). The reaction was started by the addition of 5 units DNA polymerase I (Klenow fragment). After 30 min at 37°C, the reaction was stopped by heating to 75°C (10 min).

2.22 Transfection of COS cells and CHO cells

5-10 μ g linearized plasmid DNA are dissolved in 150 μ l cell culture medium (without FKS and antibiotics). 30 μ l of the transfection agent (SuperFect, Qiagen, Hilden) are added to the solution. The mixture is incubated for 10 min at RT and added to the cells after the addition of 1 ml medium. The

COS cells (80% confluent growth) were cultivated in 60 mm plates and washed with PBS shortly before the transfection. After an incubation of 3 h at 37°C and 5% CO₂, the transfection agent was removed. The cells were then washed 4 times with PBS and taken up in selection medium (DMEM, 5% FKS, 600 µg/ml Geneticin).

2.23 Chromatographic purification of the GP120 mixture

The chromatographic purification takes place according to standard methods (Techniques in HIV Research, Aldovini & Walker, Stockton Press, 1990; S.W. Pyle et al. Purification of 120,00 dalton envelope glycoprotein from culture fluids of human immunodeficiency virus (HIV)-infected H9 cells. AIDS Res Hum Retroviruses 1987, 3:387-400). Cell extracts and cell culture overhangs of GP120-expressing COS cells are used for the isolation of the GP120 mixture. After centrifuging (25,000 g), the lysate is purified as follows:

1. Gel filtration (Sephadex G-20)
2. Immunoaffinity chromatography (anti-GP120 antibodies bound to CH-sepharose 4B)
3. Concentration through protein precipitation
4. Concentration through dialysis

2.24 Preparation of the DNA vaccine

Starting material for the preparation of the DNA vaccine is the mixture of the BstEII-BamHI DNA fragments of the env gene which were used for the preparation of the GP120 protein mixture. A eukaryotic expression vector for the HIV-1 GP120 which is approved for DNA vaccinations (cf.

e.g. J.D. Boyer et al., J Infect Dis 1997, 176:1501-1509; J. D. Boyer et al., Nat Med 1997, 3:526-532; M.L. Bagarazzi et al., J Med Primatol 1997, 26:27-33) is modified such that in the env gene the cleavage sites for BstEII and BamHI are located at identical points of the *gp120* reading frame. The mutated, variable env-gene fragments (BstEII-BamHI) are cloned into such a vector. The modification of the env genes in the V2 loop and V3 loop regions and their cloning into the DNA vaccine vector takes place analogously to the cloning steps which are carried out for the preparation of the *gp120* mixture.

"000000" 000000

Patent Claims

1. Protein vaccine which comprises a mixture of viral protein molecules, **characterized in that** the molecules are sequence variants of a single viral protein or of part of same, the mixture containing $\geq 10^2$ sequence variants, which is obtainable by expression of a plasmid-DNA mixture which, because of the variation of nucleotide positions, contains randomly distributed sequence combinations.
2. Protein vaccine according to claim 1, **characterized in that** the mixture contains $\geq 10^3$ and preferably $\geq 10^4$ sequence variants.
3. Protein vaccine according to claim 1 or 2, **characterized in that** it comprises a mixture of GP120 proteins of HIV which in each case differ from each other in their amino acid sequence in the region of the V2 loop and/or of the V3 loop.
4. DNA vaccine which codes for a mixture of structurally different virus proteins, **characterized in that** the vaccine contains a mixture of sequence variants of a viral DNA molecule or of part of same, the mixture containing $\geq 10^2$ DNA molecules which differ from each other in their nucleic acid sequence, where the mixture, because of the variation of nucleotide positions, contains randomly distributed sequence combinations.

5. DNA vaccine according to claim 4, **characterized in that** it contains a mixture of DNA molecules which code for the sequence variants of a viral protein or a part.
6. DNA vaccine according to claim 4 or 5, **characterized in that** the mixture contains $\geq 10^3$ and preferably $\geq 10^4$ DNA molecules which differ from each other in their nucleic acid sequence.
7. DNA vaccine according to one of claims 4 to 6, **characterized in that** it codes for a mixture of structurally different GP120 proteins of HIV, in which the vaccine contains a mixture of DNA molecules, the nucleic acid sequences of which differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop.
8. DNA vaccine according to claim 7, **characterized in that** it contains a mixture of DNA molecules which differ from each other in their nucleic acid sequence such that they code for a mixture of GP120 proteins which contain amino acid sequences which differ from each other in the V2 loop and/or in the V3 loop.
9. Nucleic acid sequence which is derived from the env sequence represented in SEQ ID NO: 1 or a fragment of same, **characterized in that** it is modified such that it contains monovalent restriction sites which make possible the specific exchange of the V2 and V3 regions.

10. Nucleic acid sequence which is derived from the env sequence represented in SEQ ID NO: 1 or a fragment of same, **characterized in that** it is modified such that it contains ten monovalent restriction cleavage sites at intervals of approx. 150 base pairs.
11. Nucleic acid sequence according to claim 9 or 10, **characterized in that** the sequence is modified by the introduction of silent mutations.
12. Nucleic acid sequence according to claims 9 to 11, **characterized in that** it contains the sequence given in SEQ ID NO: 9.
13. Nucleic acid sequence, **characterized in that** it contains the sequence given in SEQ ID NO: 11.
14. Nucleic acid sequence, **characterized in that** it contains the sequence given in SEQ ID NO: 12.
15. Single-stranded nucleic acid sequence which contains the region coding for the V3 loop and/or for the V2 loop of GP120, **characterized in that** in the V3 loop a 247 bp-long BglII-XbaI fragment or a 283 bp-long BglII-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-BclI fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions.

16. Single-stranded nucleic acid sequence which contains the region coding for the V2 loop and/or the region of GP120 coding for the V3 loop, **characterized in that** in the V3 loop a 247 bp-long BglIII-XbaI fragment or a 283 bp-long BglIII-NheI fragment is exchanged for a modified fragment, and in the V2 loop a 139 bp-long PstI-BclI fragment or a 339 bp-long PstI-EcoRI fragment is exchanged for a modified fragment, the fragment/the fragments in each case containing inosine, a nucleic acid exchange or a mutation at at least 6 positions, the nucleic acid sequence being complementary to the nucleic acid sequence according to claim 14.
17. Nucleic acid sequence according to claim 15 or 16, characterized in that the fragment/the fragments contain(s) inosine, a nucleic acid exchange or a mutation at 9 to 20 positions.
18. Double-stranded DNA which comprises hybrids of the single-stranded nucleic acid sequence according to claim 15 or 17 with the single-stranded nucleic acid sequence according to claim 16 or 17.
19. Nucleic acid mixture which comprises double-stranded DNAs, the nucleic acid sequences of which are derived from the env sequence in SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same, **characterized in that** the nucleic acid sequences in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop.

20. Nucleic acid mixture according to claim 19,
characterized in that the nucleic acid sequences
differ from each other such that they code for a
mixture of proteins which in each case contain amino
acid sequences which differ from each other in the V2
loop and/or in the V3 loop.
21. Nucleic acid mixture according to claim 20,
characterized in that the mixture contains $\geq 10^2$
sequence variants.
22. Nucleic acid sequence according to claim 21,
characterized in that the mixture contains $\geq 10^3$ and
preferably $\geq 10^4$ sequence variants.
23. Protein mixture which comprises sequence variants of
the GP120 protein, **characterized in that** it is a
mixture of proteins which contain amino acid sequences
which in each case differ from each other in the V2
loop and/or in the V3 loop, the mixture being
obtainable by expression of a plasmid DNA mixture
which, because of the variation of nucleotide
positions, contains randomly distributed sequence
combinations.
24. Protein mixture according to claim 23, **characterized**
in that the mixture contains $\geq 10^2$ sequence variants.
25. Protein mixture according to claim 24, **characterized**
in that the mixture contains $\geq 10^3$ and preferably $\geq 10^4$
sequence variants.

26. Plasmid, which contains an inserted double-stranded DNA according to claim 18.
27. Expression vector, characterized in that it contains an inserted nucleic acid sequence according to claims 9 to 14.
28. Expression vector according to claim 27, characterized in that it contains the sequence given in SEQ ID NO: 10.
29. Expression vector, characterized in that it corresponds to DSM 12612.
30. Vector mixture which contains a mixture of plasmids according to claim 26, characterized in that the nucleic acid sequences of the plasmids differ in each case from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop, where the mixture of the plasmids, because of the variation of nucleotide positions, contains randomly distributed sequence combinations.
31. Vector mixture according to claim 30, characterized in that the mixture contains $\geq 10^2$ plasmids which differ from each other in their nucleic acid sequence.
32. Vector mixture according to claim 31, characterized in that the mixture contains $\geq 10^3$ and preferably $\geq 10^4$

plasmids which differ from each other in their nucleic acid sequence.

33. Vector mixture according to one of claims 30 to 32, **characterized in that** the plasmids can be expressed in *E. coli* as host cell.
34. Vector mixture according to one of claims 30 to 32, **characterized in that** the plasmids can be expressed in eukaryotic cells, preferably in Cos, CHO or BHK cells, as host cells.
35. *E. coli* host cells which are transfected with a vector mixture according to claim 33.
36. Eukaryotic host cells which are transfected with a vector mixture according to claim 34.
37. Eukaryotic host cell according to claim 36, **characterized in that** they are a host cell from the group consisting of Cos, BHK or CHO cells.
38. Process for the preparation of the nucleic acid sequence according to claim 10, **characterized in that** so many silent mutations are introduced into a nucleic acid sequence coding for a viral protein that the nucleic acid sequence thereby obtained contains monovalent restriction sites which make possible the exchange of the V2 and V3 regions.
39. Process for the preparation of the nucleic acid sequence according to claim 10, **characterized in that**

so many silent mutations are introduced into a nucleic acid sequence coding for a viral protein that the nucleic acid sequence thereby obtained contains ten monovalent restriction sites at intervals of approx. 150 base pairs.

40. Process according to claim 38 or 39, **characterized in that** the nucleic acid sequence coding for a viral protein is the sequence according SEQ ID NO: 1 or SEQ ID NO: 11 or a fragment of same.
41. Process for the preparation of the vector mixture according to claims 33 and 34, **characterized in that** plasmids, the nucleic acid sequences of which in each case differ from each other in the region coding for the V2 loop and/or in the region coding for the V3 loop in each case through random distribution of the bases at the varied nucleotide positions, are ligated into a vector which can be expressed in host cells.
42. Process according to claim 41, characterized in that the host cells are *E. coli*, Cos, CHO or BHK cells.
43. Process for the preparation of the host cells according to claim 35 or 36, **characterized in that** the host cells are transformed with a vector mixture according to claims 30 to 32.
44. Process for the preparation of a protein vaccine according to one of claims 1 to 3, **characterized in that** the host cells are cultivated according to one of claims 35 to 37 under conditions which allow the

expression of the mixture of viral protein sequence variants.

45. Process for the preparation of a DNA vaccine according to one of claims 4 to 8, **characterized in that** the process is carried out according to claim 41 or 42, the plasmids according to the invention being ligated into a vector which can be expressed in host cells of the organism to be vaccinated.
46. Use of a mixture of structurally different viral proteins which are sequence variants of a viral protein or of part of same, for the preparation of a vaccine for the prevention and/or therapy of a virus infection in humans.
47. Use of a protein mixture according to one of claims 23 to 25 for the preparation of a vaccine for the prevention and/or therapy of a HIV infection in humans.
48. Use of a mixture of DNA molecules which code for sequence variants of a viral protein or of part of same, for the preparation of a vaccine for the prevention and/or therapy of a virus infection in humans.
49. Use of a nucleic acid mixture according to claims 19 to 22 for the preparation of a vaccine for the prevention and/or therapy of a virus infection in humans.

50. Use of the nucleic acid mixture according to one of claims 19 to 22 for the preparation of a vector mixture according to one of claims 30 to 32 which can be expressed in host cells , the host cells being selected from the group consisting of *E. coli*, Cos, CHO and BHK cells.
51. Use of the vector mixture according to one of claims 30 to 32 for the expression of a protein mixture according to one of claims 23 to 25.
52. Use of the host cell according to one of claims 35 to 37 for the preparation of a protein mixture according to one of claims 23 to 25.
53. Pharmaceutical composition for the prevention and/or therapy of a virus infection, **characterized in that** it comprises a protein mixture and a nucleic acid mixture, the protein mixture comprising sequence variants of a viral protein or of part of same, and the nucleic acid mixture comprising DNA molecules which code for sequence variants of a viral protein or of part of same.
54. Pharmaceutical composition according to claim 53, **characterized in that** it comprises a protein mixture according to claims 23 to 25 and a nucleic acid mixture according to claims 19 to 22.

Abstract

The present invention relates to a pharmaceutical composition or a vaccine which comprises a mixture of viral protein molecules which are sequence variants of a single viral protein or of part of same. The invention furthermore relates inter alia to a DNA vaccine which codes for a mixture of structurally different virus proteins, the vaccine containing a mixture of sequence variants of a viral DNA molecule or of part of same which code for sequence variants of a viral protein or part. According to a preferred version of the invention, the viral proteins are sequence variants of the GP120 protein of the human immunodeficiency virus (HIV) which differ from each another in their amino acid sequence in the region of the V2 loop and/or of the V3 loop. The invention furthermore relates to the preparation of the virus vaccines including the intermediate stages or constructs, preparation processes and uses connected with them.

T00130" 627600

Fig. 1

V3 loop sequence data of HIV-1 patient isolates (PI)

Fig. 2a

Amino acid sequences of the NL4-3 V2 loop and NL4-3 V3 loop. The regions which are preferably to be varied are underlined.

Fig. 2b

Variation of the V3 loop. Data from the Los Alamos database.

Fig. 4

Schematic representation of the process for the preparation of the mixture of gp 120-expressing plasmid vectors.

- 1.) Preparation of the degenerated DNA fragments for e.g. the V3 loop
 - a.) Synthesis of single-stranded DNA
 - b.) Hybridization of two complementary oligonucleotides.
- 2.) Cloning of the V3 loop DNA fragments into pUC18 delta env
- 3.) Cloning of the env gene into the gp 120 expression vector pBSCenvATG

V3 loop fragments with degenerated sequence e.g.:

Fig. 5

Heterogeneity of the vaccine using the V3 loop as an example

V3 loop sequence:

Degeneration at protein level:

variants

Degenerated DNA sequence of the respective variable amino acid positions:

variants

Variability at protein level derived from the degenerated DNA sequence:

RULE 63 (37 C.F.R. 1.63)
INVENTORS DECLARATION FOR PATENT APPLICATION
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

As a below named inventor, I hereby declare that my residence, mailing address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

VIRAL VACCINE

the specification of which (check applicable box(es)).

☐ is attached hereto
☐ was filed on _____ as U.S. Application Serial No. _____ (Atty Dkt. No. 35-213)
☒ was filed as PCT International application No. PCT/EP99/09759 on 3 December 1999
and (if applicable to U.S. or PCT application) was amended on 2 November 2000

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose to the Patent Office all information known to me to be material to patentability as defined in 37 C.F.R. 1.56. I hereby claim foreign priority benefits under 35 U.S.C. 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed or, if no priority is claimed, before the filing date of this application:

Priority Foreign Application(s):

Application Number	Country	Day/Month/Year Filed
199 07 485.2	GERMANY	12 February 1999

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

Application Number	Date/Month/Year Filed
--------------------	-----------------------

I hereby claim the benefit under 35 U.S.C. 120/365 of all prior United States and PCT international applications listed above or below.


Prior U.S./PCT Application(s):

Application Serial No.	Day/Month/Year Filed	Status: patented pending, abandoned
PCT/EP99/09759	3 December 1999	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon. And on behalf of the owner(s) hereof, I hereby appoint **NIXON & VANDERHYE P.C., 1100 North Glebe Rd., 8th Floor, Arlington, VA 22201-4714, telephone number (703) 816-4000 (to whom all communications are to be directed)**, and the following attorneys thereof (of the same address) individually and collectively owner's/owners' attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent: Larry S. Nixon, 25640; Arthur R. Crawford, 25327; James T. Hosmer, 30184; Robert W. Faris, 31352; Richard G. Besha, 22770; Mark E. Nusbaum, 32348; Michael J. Keenan, 32106; Bryan H. Davidson, 30251; Stanley C. Spooner, 27393; Leonard C. Mitchard, 29009; Duane M. Byers, 33363; Jeffrey H. Nelson, 30481; John R. Lastova, 33149; H. Warren Burnam, Jr., 29366; Mary J. Wilson, 32955; J. Scott Davidson, 33489; Alan M. Kagen, 36178; Robert A. Molan, 29834; B. J. Sadoff, 36663; James D. Berquist, 34776; Updeep S. Gill, 37334; Michael J. Shea, 34725; Donald L. Jackson, 41090; Michelle N. Lester, 32331; Frank P. Presta, 19828; Joseph S. Presta, 35329; Joseph A. Rhoa, 37515; Raymond Y. Mah, 41426; Chris Comuntzis, 31097. I also authorize Nixon & Vanderhye to delete any attorney names/numbers no longer with the firm and to act and rely solely on instructions directly communicated from the person, assignee, attorney, firm, or other organization sending instructions to Nixon & Vanderhye on behalf of the owner(s).

1. Inventor's Signature: [Signature] Date: 10-29-2001
Inventor: Michael SCHREIBER Germany
(first) (last) (citizenship)
Residence: (city) Hamburg MI (state/country) Germany
Mailing Address: Heidelberg 35, Hamburg, Germany
(Zip Code) D-22301

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☒ Page(s) _____ of DRAWINGS were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ *Scanned copy is best available.*

SCANNED, # 14